

AECL-9176

**ATOMIC ENERGY
OF CANADA LIMITED**



**L'ÉNERGIE ATOMIQUE
DU CANADA LIMITÉE**

**TABLES OF TENSOR FEYNMAN INTEGRALS
IN THE LIGHT-CONE GAUGE WITH THE
MANDELSTAM-LEIBBRANDT PRESCRIPTION**

**TABLEAUX D'INTÉGRALES TENSORIELLES DE
FEYNMAN POUR LES THÉORIES DE LA JAUGE À CÔNE LUMINEUX
DE LA PRESCRIPTION DE MANDELSTAM-LEIBBRANDT**

M.S. MILGRAM and H.C. LEE

Chalk River Nuclear Laboratories

Laboratoires nucléaires de Chalk River

Chalk River, Ontario

March 1987 mars

ATOMIC ENERGY OF CANADA LIMITED

TABLES OF TENSOR FEYNMAN INTEGRALS IN THE LIGHT-CONE GAUGE
WITH THE MANDELSTAM-LEIBBRANDT PRESCRIPTION

by

M.S. Milgram[†] and H.C. Lee^{*}

Reactor Physics Branch[†] and Theoretical Physics Branch^{*}
Chalk River Nuclear Laboratories
Chalk River, Ontario, Canada K0J 1J0
1987 March

AECL-9176

L'ÉNERGIE ATOMIQUE DU CANADA, LIMITÉE


TABLEAUX D'INTÉGRALES TENSORIELLES DE FEYNMAN POUR LES THÉORIES DE LA
JAUGE À CÔNE LUMINEUX DE LA PRESCRIPTION DE MANDELSTAM-LEIBBRANDT

par

M.S. Milgram⁺ et H.C. Lee*

RÉSUMÉ

On évalue et présente sous forme de tableaux une série complète d'intégrales tensorielles à deux points ayant jusqu'à 3 indices de Lorentz pour les théories de la jauge à cône lumineux définies dans la prescription de Mandelstam-Leibbrandt.



Physique des réacteurs⁺ et Physique des théories*
Laboratoires Nucléaires de Chalk River
Chalk River, Ontario, Canada K0J 1J0
1987 mars

AECL-9176

ATOMIC ENERGY OF CANADA LIMITED

TABLES OF TENSOR FEYNMAN INTEGRALS IN THE LIGHT-CONE GAUGE
WITH THE MANDELSTAM-LEIBBRANDT PRESCRIPTION

M.S. Milgram[†] and H.C. Lee^{*}

A b s t r a c t

A comprehensive set of two-point tensor integrals with up to 3 Lorentz indices for Yang-Mills theories in the light-cone gauge defined in the Mandelstam-Leibbrandt prescription is evaluated and presented in tabulated form.

Reactor Physics Branch[†] and Theoretical Physics Branch^{*}
Chalk River Nuclear Laboratories
Chalk River, Ontario, Canada K0J 1J0
1987 March

AECL-9176

In previous papers¹ a large set of two-point Feynman integrals encountered in one-loop computations of Yang-Mills theories in the covariant and axial gauges was evaluated using the method of analytic regularization² and presented in tabulated form. By two-point integrals we mean integrals with one external momentum that are most naturally associated with the vacuum polarization of two-point functions. Although a general description of the vacuum polarization of an n-point function calls for the evaluation of n-point integrals, it has been shown that, by judicious exploitation of symmetry, a knowledge of the relevant two-point integrals alone is sufficient to determine the one-loop counter Lagrangian which involves two-, three-, and four-point functions.³

Among the integrals given in ref. 1 are those appropriate for the light-cone gauge⁴ defined by the principal-value (PV) prescription.⁵ The light-cone gauge is the special axial gauge⁶ defined by a null vector n_+ ,

$$\underline{A} \cdot n_+ \equiv A^+ = 0, \quad n_+^2 = 0, \quad (1)$$

and the PV prescription in this case defines the inverse of the quantity $p^+ = p \cdot n_+$ as

$$\frac{1}{p^+} = \lim_{\eta \rightarrow 0^+} \frac{p^+}{(p^+)^2 + \eta} \quad (2)$$

Recently it has been shown unequivocally that the light-cone gauge prescribed by (2) is nonrenormalizable.⁷ On the other hand, it has been demonstrated that the light-cone gauge is one-loop renormalizable³ if p^+ is defined by the Mandelstam-Leibbrandt (ML) prescription⁸ ($p^- \equiv p \cdot n_-$, where n_- is the null vector conjugate to n_+)

$$\frac{1}{p^+} = \lim_{\eta \rightarrow 0^+} \frac{1}{p^+ + i\eta p^-} \quad \text{or} \quad \lim_{\eta \rightarrow 0^+} \frac{p^-}{p^+ p^- + i\eta} \quad (3)$$

In this paper we give the values of a comprehensive set of two-point tensor light-cone gauge integrals based on (3). These integrals have become especially relevant following the recent realization⁹ that superstring theories can be easily quantized only in the light-cone gauge.

As in ref. 1, the scalar integrals are evaluated via an analytic representation for the generalized two-point integrals

$$\begin{aligned} M(\omega, \kappa, \mu, \nu, \lambda; p) &\equiv \int d^2 q \left[(p-q)^2 \right]^\kappa (q^2)^\mu (q^+)^{\nu} (q^-)^{\lambda} \\ &= \frac{i(-\pi)^\omega (p^2)^{\alpha_1 - \nu} (p^+)^{\nu} (p^-)^{\lambda} \Gamma(\lambda+1)}{\Gamma(-\kappa) \Gamma(-\mu) \Gamma(-\nu) \Gamma(2\omega + \kappa + \mu + \nu + \lambda)} \\ &\quad \times u^{-\nu} G_{3,3}^{2,3} \left(u \left| \begin{matrix} 1 + \nu, 1 - \omega - \mu - \lambda, 1 + \alpha_1; \\ 0, \omega + \kappa + \nu; \nu - \lambda \end{matrix} \right. \right), \end{aligned} \quad (4)$$

$$\alpha_1 = \omega + \kappa + \mu + \nu, \quad u = 2p^+ p^- / p^2$$

where κ , μ , ν and λ are continuous exponents, 2ω is the continuous dimension of spacetime¹⁰ and G is a Meijer G-function.¹¹ The derivation of (4) and its various simplifying special cases are given in detail elsewhere. When ν is a nonnegative integer and $\lambda = 0$, (4) reduces to a representation identical to that for the PV prescription. When $\nu < 0$ the two prescriptions give different results. In particular the ML prescription obeys the rule of power counting whereas the PV prescription does not.

The integral (4) is ultraviolet (UV) divergent when the UV index α_1 satisfies

$$\alpha_1 = \omega + \kappa + \mu + \nu = \text{integer} \geq 0 \quad (5)$$

and is infrared (IR) divergent when either one or both of the conditions

$$\omega + \mu + \lambda = \text{integer} \leq 0 \quad (6)$$

$$\omega + \kappa + \nu = \text{integer} \leq 0$$

is or are met. The limiting process

$$\begin{aligned} \nu, \lambda &= (N, L) \text{ integers, } \kappa, \mu = (K, M) \text{ integers} + \sigma, \quad \omega = 2 + \varepsilon \\ \sigma &\rightarrow 0, \quad \varepsilon \rightarrow \text{small} \end{aligned} \quad (7)$$

separates the UV and IR poles in (4) and preserves gauge invariance.¹² In the tabulated results UV poles are of $O(1/e_1)$ and IR poles of $O(1/e_0)$, where in the limit (7) e_0 and e_1 have identical values, viz.

$$\frac{1}{e_n} = \frac{1}{\varepsilon + 2\sigma - n\sigma} + \gamma + \lambda n p^2 \rightarrow \frac{1}{\varepsilon} + \gamma + \lambda n p^2, \quad n = 0, 1 \quad (8)$$

Because in studies of quantum field theories the UV divergent part of vertex functions are usually of particular interest, the integrals to be tabulated will be classified according to the UV index, taking on all values within the range

$$-2 \leq \alpha_1 \leq 2 \quad (9)$$

With this classification, integrals have been evaluated for the following ranges of the (integer) exponents

$$\begin{aligned} -2 &\leq K \leq -1 \\ -2 &\leq M \leq 2 \\ -2 &\leq N \leq 2 \\ 0 &\leq L \leq 2 \end{aligned} \quad (10)$$

We remark that in the light-cone gauge integrals with $\lambda < 0$ never occur. Note also that integrals with $\lambda = 0$ and $\nu > 0$ are already given in ref. 1, but for completeness they will be given again here. So called "tadpole" integrals satisfying either or both of the following conditions

$$\begin{aligned} \kappa &\geq 0 ; \\ \mu &\geq 0 \quad \text{and} \quad \nu \geq 0 \end{aligned} \tag{11}$$

vanish in the limit (7).

We now note that although (4) only applies to integrals with integrands not containing uncontracted Lorentz indices, any integral with indices can be straightforwardly reduced³ to a linear combination of those without indices. It was shown in ref. 3 that, for tensors of up to rank-4, the reduction is achieved by replacing tensor integrands as follows:

$$\begin{aligned} q_i q_j q_k q_l &\rightarrow \frac{1}{3} (x-y^2) p^4 (\delta_{ij} \delta_{kl} + 2 \text{ symmetric terms}) \\ &- \frac{1}{3} (x-y^2)(x-4y^2) p^2 (p_i p_j \delta_{kl} + 5 \text{ terms}) \\ &+ (x^2 - 8xy^2 + 8y^4) p_i p_j p_k p_l \end{aligned} \tag{12a}$$

$$q_i q_j q_k \rightarrow y(x-y^2) p^2 (\delta_{ij} p_k + 2 \text{ terms}) + y(-3x+4y^2) p_i p_j p_k \tag{12b}$$

$$q_i q_j \rightarrow (x-y^2) p^2 \delta_{ij} + (-x+2y^2) p_i p_j \tag{12c}$$

$$q_i \rightarrow y p_i \tag{12d}$$

where

$$p^2 \equiv (1-u) p^2 \tag{13a}$$

$$p^2_x \equiv q^2 = 2q^+ q^- - q^2 \tag{13b}$$

$$p^2_y \equiv \dot{p} \cdot q = p^+ q^- + p^- q^+ - \frac{1}{2} [p^2 + q^2 - (p-q)^2] \tag{13c}$$

In conjunction with these relations, the expressions (4) and (12) suffice to cover all two-point integrals (up to rank-4) that will ever need to be evaluated in the light-cone gauge. The ultraviolet condition (5) becomes

$$\omega + \kappa + \mu + \nu + R = \alpha_1 + R \geq 0 \tag{5'}$$

and the infrared conditions (6) become

$$\begin{aligned} \omega + \mu + \lambda + R &\leq 0 \\ \omega + \kappa + \nu + R &\leq 0 \end{aligned} \tag{6'}$$

where R is the rank of the tensor integrand. Integrals with integrands containing one-, two- and three-component tensors have been computed using (12) and tabulated for the same range of variables cited in (9) and (10) in the accompanying Tables 1-8. The computer code used to evaluate the integrals is completely general for any value of the indices $\kappa, \mu, \nu, \lambda > 0$ (and ω if need be), and tensor integrands with up to 4 uncontracted Lorentz indices.

The integrals in the Tables for values of α_1 running between -2 and 2 , are labelled by their individual exponents. The integrals are normalized by the omission of a factor

$$f = i(-\pi)^\omega (p^2)^{\alpha_1 - N} (p^+)^N (p^-)^L \quad (12)$$

so that with the exception of the two quantities $1/\hat{e}_0$ and $1/\hat{e}_1$, they are functions of the single variable $u = 2p^+p^-/p^2$. The finite parts sometimes contain an infinite sum defined by

$$S_n = \sum_{\ell=0}^{\infty} \frac{u^{\ell+1}}{\ell+n} \left(\log u - \frac{1}{\ell+n} \right) \quad (13)$$

The generation of the Tables from the representation (4) was carried out on CDC computers using the algebraic programming code SCHOONSCHIP¹³, and on-line editors.

References

1. M.S. Milgram and H.C. Lee, J. Comp. Phys. 59, 331 (1985); J. Comp. Phys. (in press for Vol. 71, no. 2, Aug. 1987).
2. H.C. Lee and M.S. Milgram, Phys. Lett. 133B, 320 (1983); Ann. Phys. 157, 408 (1984).
3. H.C. Lee and M.S. Milgram, Phys. Rev. Lett. 55, 2122 (1985) and Nucl. Phys. B 268, 543 (1986), where it is shown that although the result for individual proper vertex functions at one-loop order in the light-cone gauge may suggest the need for nonlocal counterterms, all nonlocal counterterms cancel exactly in the total counter-Lagrangian.
4. J.M. Cornwall, Phys. Rev. D10(1974)500; D26, 1453 (1982); L. Brink, O. Lindgren and B.E.W. Nilsson, Nucl. Phys. B212, 401 (1983); S. Mandelstam, Nucl. Phys. B213(1983)149; G. Leibbrandt, Phys. Rev. D29, 1699 (1984); G. Leibbrandt and S.L. Nyeo, Phys. Lett. 140B, 417 (1984); D.M. Capper, J.J. Dulwich and M.J. Litvak, Nucl. Phys. B241 463 (1984); H.C. Lee and M.S. Milgram, Refs. 2 and 3.
5. W. Kummer, Acta. Phys. Austr. 41, 315 (1975)
6. W. Kummer, Acta. Phys. Austr. 15, 149 (1961); A. Salam and J. Strathdee, Nuovo Cimento 23A, 237 (1974); D.M. Capper and G. Leibbrandt, Phys. Rev. D25, 1002,1009 (1982); H.C. Lee and M.S. Milgram, ref. 2 and J. Math. Phys. 26, 1793 (1965).
7. Cornwall, ref. 4 (1982); Capper, Dulwich and Litvak, ref. 4; G. Leibbrandt and T. Matsuki, Phys. Rev. D31, 934 (1985); H.C. Lee and M.S. Milgram, Z. Phys. C28, 579 (1985); H.C. Lee, Chinese J. Phys. 23, 90 (1985).
8. S. Mandelstam, ref. 4; G. Leibbrandt, ref. 4.
9. M.B. Green and J.H. Schwarz, Nucl. Phys. B243, 285,475 (1984).
10. G. 'tHooft and M. Veltman, Nucl. Phys. B44, 189 (1972); C.G. Bellini and J.J. Giambiagi, Nuova Cimento B12, 20 (1972); J.F. Ashmore, Nuovo Cimento Lett. 4, 289 (1972).
11. Y. Luke, "The Special Functions and Their Approximations", Vol. 1, Chap. 5 (Academic, New York, 1969).
12. H.C. Lee and M.S. Milgram, ref. 6.
13. M. Veltman, Univ. of Michigan (1964; revised (1984); H. Strubbe, Comp. Phys. Comm. 18, 1 (1979).

TABLE 1(A) $K=-2, \alpha_1=-2$

$$[K, M, N, L] = f \int d^4 q ((p-q)^2)^K (q^2)^{2M} (q^+)^N (q^-)^L$$

$$\begin{aligned} [-2, 0, -2, 0] &= -1 + \log u + 1/\hat{e}_0 \\ [-2, -1, -1, 0] &= \log u + 1/\hat{e}_0 \\ [-2, -2, 0, 0] &= -2 + 2/\hat{e}_0 \\ [-2, 0, -2, 1] &= -2 + \log u + 1/\hat{e}_0 \\ [-2, -1, -1, 1] &= -1 + \log u + 1/\hat{e}_0 + s_2 \\ [-2, -2, 0, 1] &= -1 + 1/\hat{e}_0 \\ [-2, 0, -2, 2] &= -5/2 + \log u + 1/\hat{e}_0 \\ [-2, -1, -1, 2] &= -3/2 + \log u + 1/\hat{e}_0 + 2s_3 \\ [-2, -2, 0, 2] &= -2 + 1/\hat{e}_0 \end{aligned}$$

TABLE 1(B) $K=-2, \alpha_1=-1$

$$[K, M, N, L] = f \int d^4 q ((p-q)^2)^K (q^2)^{2M} (q^+)^N (q^-)^L$$

$$\begin{aligned} [-2, 1, -2, 0] &= -1 + \log u + u + 1/\hat{e}_0 \\ [-2, 0, -1, 0] &= \log u + 1/\hat{e}_0 \\ [-2, -1, 0, 0] &= 1/\hat{e}_0 \\ [-2, -2, 1, 0] &= -1 + 1/\hat{e}_0 \\ [-2, 1, -2, 1] &= -2 + \log u + u + 1/\hat{e}_0 \\ [-2, 0, -1, 1] &= -1 + \log u + 1/\hat{e}_0 \\ [-2, 1, -2, 2] &= -1 + 1/\hat{e}_0 \\ [-2, 0, -1, 2] &= -3/2 + \log u + 1/\hat{e}_0 \\ [-2, -1, 0, 2] &= -3/2 + 1/\hat{e}_0 \\ [-2, -2, 1, 2] &= -5/2 + 1/u + 1/\hat{e}_0 \end{aligned}$$

TABLE 1(C) K=-2, $\alpha_1=0$

$$[K, M, N, L] = f \int d^4 q \left((p-q)^2 \right)^K (q^+)^2 M (q^+)^N (q^-)^L$$

$[-2, 2, -2, 0]$	$= -1 + \log u - 3u^2 \log u + 4u - 7u^2/2 + 1/\hat{e}_0 + 1/\hat{e}_1 * (-3u^2)$
$[-2, 1, -1, 0]$	$= \log u - 2u \log u + 1/\hat{e}_0 + 1/\hat{e}_1 * (-2u)$
$[-2, 0, 0, 0]$	$= 1/\hat{e}_0 - 1/\hat{e}_1$
$[-2, -1, 1, 0]$	$= -1 + 1/\hat{e}_0$
$[-2, -2, 2, 0]$	$= -2 + 1/\hat{e}_0$
$[-2, 2, -2, 1]$	$= -2 + \log u - 2u^2 \log u + 3u - 2u^2/3 + 1/\hat{e}_0 + 1/\hat{e}_1 * (-2u^2)$
$[-2, 1, -1, 1]$	$= -1 + \log u - 3u/2 \log u + 5u/4 + 1/\hat{e}_0 + 1/\hat{e}_1 * (-3u/2)$
$[-2, 0, 0, 1]$	$= 1/\hat{e}_0 - 1/\hat{e}_1$
$[-2, -1, 1, 1]$	$= -3/2 + 1/u + 1/\hat{e}_0 + 1/\hat{e}_1 * (-1/(2u))$
$[-2, -2, 2, 1]$	$= -5/2 + 1/u + 1/\hat{e}_0$
$[-2, 2, -2, 2]$	$= -5/2 + \log u - 5u^2/3 \log u + 8u/3 + 11u^2/36 + 1/\hat{e}_0 + 1/\hat{e}_1 * (-5u^2/3)$
$[-2, 1, -1, 2]$	$= -3/2 + \log u - 4u/3 \log u + 16u/9 + 1/\hat{e}_0 + 1/\hat{e}_1 * (-4u/3)$
$[-2, 0, 0, 2]$	$= 1/\hat{e}_0 - 1/\hat{e}_1$
$[-2, -1, 1, 2]$	$= -11/6 + 13/(9u) + 1/\hat{e}_0 + 1/\hat{e}_1 * (-2/(3u))$
$[-2, -2, 2, 2]$	$= -17/6 + 5/(9u^2) + 4/(3u) + 1/\hat{e}_0 + 1/\hat{e}_1 * (-1/(3u^2))$

TABLE 1(D) K=-2, $\alpha_1 = 1$

$$\{K, M, N, L\} = f \int d^4 q ((p-q)^2)^K (q^2)^M (q^+)^N (q^-)^L$$

$[-2, 2, -1, 0]$	$= \log u - 6u \log u + 6u^2 \log u + 2u - 2u^2 + 1/e_0 + 1/e_1 * (-6u + 6u^2)$
$[-2, 1, 0, 0]$	$= 1/e_0 - 1/e_1$
$[-2, 0, 1, 0]$	$= 1/e_0 - 1/e_1$
$[-2, -1, 2, 0]$	$= -3/2 + 1/e_0$
$[-2, 2, -1, 1]$	$= -1 + \log u - 4u \log u + 10u^2/3 \log u + 4u - 28u^2/9 + 1/e_0 + 1/e_1 * (-4u + 10u^2/3)$
$[-2, 1, 0, 1]$	$= 1/e_0 - 1/e_1$
$[-2, 0, 1, 1]$	$= 1/e_0 - 1/e_1$
$[-2, -1, 2, 1]$	$= -11/6 + 13/(9u) + 1/e_0 + 1/e_1 * (-2/(3u))$
$[-2, 2, -1, 2]$	$= -3/2 + \log u - 10u/3 \log u + 5u^2/2 \log u + 43u/9 - 27u^2/8 + 1/e_0 + 1/e_1 * (-10u/3 + 5u^2/2)$
$[-2, 1, 0, 2]$	$= 1/e_0 - 1/e_1$
$[-2, 0, 1, 2]$	$= 1/e_0 - 1/e_1$
$[-2, -1, 2, 2]$	$= -25/12 - 4/(9u^2) + 7/(3u) + 1/e_0 + 1/e_1 * (1/(6u^2) - 1/u)$

TABLE 1(E) K=-2, $\alpha_1 = 2$

$[-2, 2, 0, 0]$	$= 1/e_0 - 1/e_1$	$[-2, 2, 0, 1]$	$= 1/e_0 - 1/e_1$	$[-2, 2, 0, 2]$	$= 1/e_0 - 1/e_1$
$[-2, 1, 1, 0]$	$= 1/e_0 - 1/e_1$	$[-2, 1, 1, 1]$	$= 1/e_0 - 1/e_1$	$[-2, 1, 1, 2]$	$= 1/e_0 - 1/e_1$
$[-2, 0, 2, 0]$	$= 1/e_0 - 1/e_1$	$[-2, 0, 2, 1]$	$= 1/e_0 - 1/e_1$	$[-2, 0, 2, 2]$	$= 1/e_0 - 1/e_1$

TABLE 2(A) $K=-1, \alpha_1=-2$

$$[K,M,N,L] = f \int d^4q ((p-q)^2)^K (q^2)^M (q^+)^N (q^-)^L$$

$$[-1,-1,-2,0] = u$$

$$[-1,-2,-1,0] = -u \log u - u + u/\hat{e}_0$$

$$[-1,-1,-2,1] = u/2 - u*(S_2-2*S_3)$$

$$[-1,-2,-1,1] = -S_2$$

$$[-1,-1,-2,2] = u/3 - u*(4*S_3 - 6*S_4)$$

$$[-1,-2,-1,2] = 2 * S_2 - 4 * S_3$$

TABLE 2(B) $K=-1, \alpha_1=-1$

$$[K,M,N,L] = f \int d^4q ((p-q)^2)^K (q^2)^M (q^+)^N (q^-)^L$$

$$[-1,0,-2,0] = u$$

$$[-1,-2,0,1] = 1$$

$$[-1,-1,-1,0] = -S_1$$

$$[-1,0,-2,2] = u/3$$

$$[-1,-2,0,0] = 1/\hat{e}_0$$

$$[-1,-1,-1,2] = -S_3$$

$$[-1,0,-2,1] = u/2$$

$$[-1,-2,0,2] = 1/2$$

$$[-1,-1,-1,1] = -S_2$$

TABLE 2(C) $K=-1, \alpha_1=0$

$$[K,M,N,L] = f \int d^4q ((p-q)^2)^K (q^2)^M (q^+)^N (q^-)^L$$

$$[-1,1,-2,0] = -u^2 \log u + u - u^2/2 + 1/\hat{e}_1 * (-u^2)$$

$$[-1,0,-1,0] = -u \log u + u + 1/\hat{e}_1 * (-u)$$

$$[-1,-1,0,0] = 2 - 1/\hat{e}_1$$

$$[-1,-2,1,0] = 1$$

$$[-1,1,-2,1] = -u^2/2 \log u + u/2 + u^2/12 + 1/\hat{e}_1 * (-u^2/2)$$

$$[-1,0,-1,1] = -u/2 \log u + 3u/4 + 1/\hat{e}_1 * (-u/2)$$

$$[-1,-1,0,1] = 1 - 1/2/\hat{e}_1$$

$$[-1,-2,1,1] = 1/2 + 1/u + 1/\hat{e}_1 * (-1/(2u))$$

$$[-1,1,-2,2] = -u^2/3 \log u + u/3 + 7u^2/36 + 1/\hat{e}_1 * (-u^2/3)$$

$$[-1,0,-1,2] = -u/3 \log u + 11u/18 + 1/\hat{e}_1 * (-u/3)$$

$$[-1,-1,0,2] = 13/18 - 1/3/\hat{e}_1$$

$$[-1,-2,1,2] = 1/3 + 5/(9u) + 1/\hat{e}_1 * (-1/(3u))$$

TABLE 2(D) K=-1, $\alpha_1=1$

$$[K,M,N,L] = f \int d^4q ((p-q)^2)^K (q^2)^M (q^+)^N (q^-)^L$$

$$[-1,2,-2,0] = -3u^2 \log u + 4u^3 \log u + u - u^2/2 - 2u^3/3 + 1/\hat{e}_1 * (-3u^2 + 4u^3)$$

$$[-1,1,-1,0] = -u \log u + 3u^2/2 \log u + u - 5u^2/4 + 1/\hat{e}_1 * (-u + 3u^2/2)$$

$$[-1,0,0,0] = 0$$

$$[-1,-1,1,0] = 1 - 1/2/\hat{e}_1$$

$$[-1,-2,2,0] = 1/2$$

$$[-1,2,-2,1] = -4u^2/3 \log u + 5u^3/3 \log u + u/2 + 4u^2/9 - 41u^3/36 + 1/\hat{e}_1 * (-4u^2/3 + 5u^3/3)$$

$$[-1,1,-1,1] = -u/2 \log u + 2u^2/3 \log u + 3u/4 - 8u^2/9 + 1/\hat{e}_1 * (-u/2 + 2u^2/3)$$

$$[-1,0,0,1] = 0$$

$$[-1,-1,1,1] = 13/18 - 4/(9u) + 1/\hat{e}_1 * (-1/3 + 1/(6u))$$

$$[-1,-2,2,1] = 1/3 + 5/(9u) + 1/\hat{e}_1 * (-1/(3u))$$

$$[-1,2,-2,2] = -5u^2/6 \log u + u^3 \log u + u/3 + 41u^2/72 - 21u^3/20 + 1/\hat{e}_1 * (-5u^2/6 + u^3)$$

$$[-1,1,-1,2] = -u/3 \log u + 5u^2/12 \log u + 11u/18 - 101u^2/144 + 1/\hat{e}_1 * (-u/3 + 5u^2/12)$$

$$[-1,0,0,2] = 0$$

$$[-1,-1,1,2] = 7/12 - 4/(9u) + 1/\hat{e}_1 * (-1/4 + 1/(6u))$$

$$[-1,-2,2,2] = 1/4 - 4/(9u^2) + 5/(9u) + 1/\hat{e}_1 * (1/(6u^2) - 1/(3u))$$

TABLE 2(E) $K=-1, \alpha_1=2$

$$[K,M,N,L] = f \int d^4q ((p-q)^2)^K (q^2)^M (q^+)^N (q^-)^L$$

$$[-1,2,-1,0] = -u \log u + 4u^2 \log u - 10u^3/3 \log u + u - 4u^2 + 28u^3/9 \\ - 1/\hat{e}_1 * (u - 4u^2 + 10u^3/3)$$

$$[-1,1,0,0] = 0$$

$$[-1,0,1,0] = 0$$

$$[-1,-1,2,0] = 13/18 - 1/3/\hat{e}_1$$

$$[-1,2,-1,1] = -u/2 \log u + 5u^2/3 \log u - 5u^3/4 \log u + 3u/4 - 43u^2/18 + 27u^3/16 \\ + 1/\hat{e}_1 * (-u/2 + 5u^2/3 - 5u^3/4)$$

$$[-1,1,0,1] = 0$$

$$[-1,0,1,1] = 0$$

$$[-1,-1,2,1] = 7/12 - 4/(9u) + 1/\hat{e}_1 * (-1/4 + 1/(6u))$$

$$[-1,2,-1,2] = -u/3 \log u + u^2 \log u - 7u^3/10 \log u + 11u/18 - 7u^2/4 + 233u^3/200 \\ + 1/\hat{e}_1 * (-u/3 + u^2 - 7u^3/10)$$

$$[-1,1,0,2] = 0$$

$$[-1,0,1,2] = 0$$

$$[-1,-1,2,2] = 149/300 + 23/(225u^2) - 41/(75u) + 1/\hat{e}_1 * (-1/5 - 1/(30u^2) + 1/(5u))$$

TABLE 3(A) $K=-2$ $\alpha_1=-2$

$$[K,M,N,L]_1 = f \int d^4q ((p-q)^2)^K (q^2)^M (q^+)^N (q^-)^L q_i$$

all terms are coefficients of p_i

$$\begin{aligned} [-2,0,-2,0]_1 &= -1 + \log u + 1/\hat{e}_0 & [-2,-2,0,1]_1 &= -2 + 1/\hat{e}_0 \\ [-2,-1,-1,0]_1 &= -1/(u-1) * \log u + 1/\hat{e}_0 & [-2,0,-2,2]_1 &= -5/2 + \log u + 1/\hat{e}_0 \\ [-2,-2,0,0]_1 &= -1 + 1/\hat{e}_0 & [-2,-1,-1,2]_1 &= -3/2 - 1/(u-1) * \log u + 1/\hat{e}_0 \\ [-2,0,-2,1]_1 &= -2 + \log u + 1/\hat{e}_0 & [-2,-2,0,2]_1 &= -5/2 + 1/\hat{e}_0 \\ [-2,-1,-1,1]_1 &= -1 - 1/(u-1) * \log u + 1/\hat{e}_0 \end{aligned}$$

TABLE 3(B) $K=-2$, $\alpha_1=-1$

$$[K,M,N,L]_1 = f \int d^4q ((p-q)^2)^K (q^2)^M (q^+)^N (q^-)^L q_i$$

all terms are coefficients of p_i

$$\begin{aligned} [-2,1,-2,0]_1 &= -1 + \log u + 2u + 1/\hat{e}_0 & [-2,-1,0,1]_1 &= -3/2 + 1/\hat{e}_0 \\ [-2,0,-1,0]_1 &= \log u + 1/\hat{e}_0 & [-2,-2,1,1]_1 &= -5/2 + 1/(2u) + 1/\hat{e}_0 \\ [-2,-1,0,0]_1 &= -1 + 1/\hat{e}_0 & [-2,1,-2,2]_1 &= -5/2 + \log u + 4u/3 + 1/\hat{e}_0 \\ [-2,-2,1,0]_1 &= -2 + 1/\hat{e}_0 & [-2,0,-1,2]_1 &= -3/2 + \log u + 1/\hat{e}_0 \\ [-2,1,-2,1]_1 &= -2 + \log u + 3u/2 + 1/\hat{e}_0 & [-2,-1,0,2]_1 &= -11/6 + 1/\hat{e}_0 \\ [-2,0,-1,1]_1 &= -1 + \log u + 1/\hat{e}_0 & [-2,-2,1,2]_1 &= -17/6 + 2/(3u) + 1/\hat{e}_0 \end{aligned}$$

TABLE 3(C) $K=-2, \alpha_1=0$

$$[K,M,N,L]_1 = \int d^4q ((p-q)^2)^K (q^2)^M (q^+)^N (q^-)^L q_i$$

all terms are coefficients of p_i

$$[-2,2,-2,0]_1 = -1 + \log u - 6u^2 \log u + 6u - 4u^2 + 1/\hat{e}_0 + 1/\hat{e}_1 * (-6u^2)$$

$$[-2,1,-1,0]_1 = \log u - 3u \log u + u + 1/\hat{e}_0 + 1/\hat{e}_1 * (-3u)$$

$$[-2,0,0,0]_1 = 1/\hat{e}_0 - 1/\hat{e}_1$$

$$[-2,-1,1,0]_1 = -3/2 + 1/\hat{e}_0$$

$$[-2,-2,2,0]_1 = -5/2 + 1/\hat{e}_0$$

$$[-2,2,-2,1]_1 = -2 + \log u - 10u^2/3 \log u + 4u - 2u^2/9 + 1/\hat{e}_0 + 1/\hat{e}_1 * (-10u^2/3)$$

$$[-2,1,-1,1]_1 = -1 + \log u - 2u \log u + 2u + 1/\hat{e}_0 - 2u/\hat{e}_1$$

$$[-2,0,0,1]_1 = 1/\hat{e}_0 - 1/\hat{e}_1$$

$$[-2,-1,1,1]_1 = -11/6 + 13/(18u) + 1/\hat{e}_0 + 1/\hat{e}_1 * (-1/(3u))$$

$$[-2,-2,2,1]_1 = -17/6 + 2/(3u) + 1/\hat{e}_0$$

$$[-2,2,-2,2]_1 = -5/2 + \log u - 5u^2/2 \log u + 10u/3 + 7u^2/8 + 1/\hat{e}_0 + 1/\hat{e}_1 * (-5u^2/2)$$

$$[-2,1,-1,2]_1 = -3/2 + \log u - 5u/3 \log u + 43u/18 + 1/\hat{e}_0 + 1/\hat{e}_1 * (-5u/3)$$

$$[-2,0,0,2]_1 = 1/\hat{e}_0 - 1/\hat{e}_1$$

$$[-2,-1,1,2]_1 = -25/12 + 7/(6u) + 1/\hat{e}_0 + 1/\hat{e}_1 * (-1/(2u))$$

$$[-2,-2,2,2]_1 = -37/12 + 5/(18u^2) + 1/u + 1/\hat{e}_0 - 1/\hat{e}_1 * (1/(6u^2))$$

TABLE 3(D) $K=-2, \alpha_1=1$

$$[K,M,N,L]_1 = f \int d^4q ((p-q)^2)^K (q^2)^M (q^+)^N (q^-)^L q_i$$

all terms are coefficients of p_i

$$\begin{aligned} [-2,2,-1,0]_1 &= \log u - 8u \log u + 10u^2 \log u + 4u - 6u^2 + 1/\hat{e}_0 + 1/\hat{e}_1 * (-8u + 10u^2) \\ [-2,1,0,0]_1 &= 1/\hat{e}_0 - 1/\hat{e}_1 \\ [-2,0,1,0]_1 &= 1/\hat{e}_0 - 1/\hat{e}_1 \\ [-2,-1,2,0]_1 &= -11/6 + 1/\hat{e}_0 \\ [-2,2,-1,1]_1 &= -1 + (-5u + 5u^2 + 1) \log u + 11u/2 - 11u^2/2 + 1/\hat{e}_0 + 1/\hat{e}_1 * (-5u + 5u^2) \\ [-2,1,0,1]_1 &= 1/\hat{e}_0 - 1/\hat{e}_1 \\ [-2,0,1,1]_1 &= 1/\hat{e}_0 - 1/\hat{e}_1 \\ [-2,-1,2,1]_1 &= -25/12 + 7/(6u) + 1/\hat{e}_0 + 1/\hat{e}_1 * (-1/(2u)) \\ [-2,2,-1,2]_1 &= -3/2 + \log u - 4u \log u + 7u^2/2 \log u + 6u - 41u^2/8 + 1/\hat{e}_0 - 1/\hat{e}_1 * (4u - 7u^2) \\ [-2,1,0,2]_1 &= 1/\hat{e}_0 - 1/\hat{e}_1 \\ [-2,0,1,2]_1 &= 1/\hat{e}_0 - 1/\hat{e}_1 \\ [-2,-1,2,2]_1 &= -137/60 - 41/(150u^2) + 149/(75u) + 1/\hat{e}_0 + 1/\hat{e}_1 * (1/(10u^2) - 4/(5u)) \end{aligned}$$

TABLE 3(E) $K=-2, \alpha_1=2$

$$[K,M,N,L]_1 = f \int d^4q ((p-q)^2)^K (q^2)^M (q^+)^N (q^-)^L q_i$$

all terms are coefficients of p_i

(This table is identical to TABLE 1(E))

TABLE 4(A) $K=-1, \alpha_1=-2$

$$[K,M,N,L]_1 = f \int d^4q ((p-q)^2)^K (q^2)^M (q^+)^N (q^-)^L q_i$$

all terms are coefficients of p_i

$$[-1,-1,-2,0]_1 = -u/(u-1)*\log u - S_1$$

$$[-1,-2,-1,0]_1 = u/(u-1)*\log u$$

$$[-1,-1,-2,1]_1 = -2 + \log u - 1/(u-1)*\log u - 2*S_1/u$$

$$[-1,-2,-1,1]_1 = 1 + 1/(u-1)*\log u + S_1/u$$

$$[-1,-1,-2,2]_1 = -3/4 + 1/2*\log u + 3/u*\log u - 1/(u-1)*\log u - 3*S_1/u^2 - 3/u$$

$$[-1,-2,-1,2]_1 = 1/2 - 2/u*\log u + 1/(u-1)*\log u + 2*S_1/u^2 + 2/u$$

TABLE 4(B) $K=-1, \alpha_1=-1$

$$[K,M,N,L]_1 = f \int d^4q ((p-q)^2)^K (q^2)^M (q^+)^N (q^-)^L q_i$$

all terms are coefficients of p_i

$$[-1,0,-2,0]_1 = u$$

$$[-1,-2,0,1]_1 = 1/2$$

$$[-1,-1,-1,0]_1 = u/(u-1)*\log u$$

$$[-1,0,-2,2]_1 = u/3$$

$$[-1,-2,0,0]_1 = 1$$

$$[-1,-1,-1,2]_1 = u/3/(u-1)*\log u$$

$$[-1,0,-2,1]_1 = u/2$$

$$[-1,-2,0,2]_1 = 1/3$$

$$[-1,-1,-1,1]_1 = u/2/(u-1)*\log u$$

TABLE 4(C) $K=-1, \alpha_1 = 0$

$$[K,M,N,L]_1 = f \int d^4q ((p-q)^2)^K (q^2)^M (q^+)^N (q^-)^L q_i$$

all terms are coefficients of p_i

$$[-1,1,-2,0]_1 = -3u^2/2 \log u + u - u^2/4 + 1/\hat{e}_1 * (-3u^2/2)$$

$$[-1,0,-1,0]_1 = -u \log u + u - u/\hat{e}_1$$

$$[-1,-1,0,0]_1 = 1 - 1/(2\hat{e}_1)$$

$$[-1,-2,1,0]_1 = 1/2$$

$$[-1,1,-2,1]_1 = -2u^2/3 \log u + u/2 + 2u^2/9 + 1/\hat{e}_1 * (-2u^2/3)$$

$$[-1,0,-1,1]_1 = -u/2 \log u + 3u/4 - 1/\hat{e}_1 * (u/2)$$

$$[-1,-1,0,1]_1 = 13/18 - 1/(3\hat{e}_1)$$

$$[-1,-2,1,1]_1 = 1/3 + 5/(18u) + 1/\hat{e}_1 * (-1/(6u))$$

$$[-1,1,-2,2]_1 = -5u^2/12 \log u + u/3 + 41u^2/144 - 1/\hat{e}_1 * (5u^2/12)$$

$$[-1,0,-1,2]_1 = -u/3 \log u + 11u/18 + 1/\hat{e}_1 * (-u/3)$$

$$[-1,-1,0,2]_1 = 7/12 - 1/(4\hat{e}_1)$$

$$[-1,-2,1,2]_1 = 1/4 + 5/(18u) + 1/\hat{e}_1 * (-1/(6u))$$

TABLE 4(D) $K=-1, \alpha_1 = 1$

$$[K,M,N,L]_1 = f \int d^4q ((p-q)^2)^K (q^2)^M (q^+)^N (q^-)^L q_i$$

all terms are coefficients of p_i

$$\begin{aligned} [-1,2,-2,0]_1 &= -4u^2 \log u + 20u^3/3 \log u + u - 26u^3/9 + 1/\hat{e}_1 * (-4u^2 + 20u^3/3) \\ [-1,1,-1,0]_1 &= u(2u-1) \log u + u - 2u^2 + 1/\hat{e}_1 * (-u + 2u^2) \\ [-1,0,0,0]_1 &= 0 \\ [-1,-1,1,0]_1 &= 13/18 - 1/(3\hat{e}_1) \\ [-1,-2,2,0]_1 &= 1/3 \\ [-1,2,-2,1]_1 &= 5u^2(u/2-1/3) \log u + u/2 + 13u^2/18 - 17u^3/8 + 1/\hat{e}_1 * (-5u^2/3 + 5u^3/2) \\ [-1,1,-1,1]_1 &= -u/2 \log u + 5u^2/6 \log u + 3u/4 - 43u^2/36 + 1/\hat{e}_1 * (-u/2 + 5u^2/6) \\ [-1,0,0,1]_1 &= 0 \\ [-1,-1,1,1]_1 &= 7/12 - 2/(9u) + 1/\hat{e}_1 * (-1/4 + 1/(12u)) \\ [-1,-2,2,1]_1 &= 1/4 + 5/(18u) + 1/\hat{e}_1 * (-1/(6u)) \\ [-1,2,-2,2]_1 &= -u^2 \log u + 7u^3/5 \log u + u/3 + 3u^2/4 - 163u^3/100 + 1/\hat{e}_1 * (-u^2 + 7u^3/5) \\ [-1,1,-1,2]_1 &= -u/3 \log u + u^2/2 \log u + 11u/18 - 7u^2/8 + 1/\hat{e}_1 * (-u/3 + u^2/2) \\ [-1,0,0,2]_1 &= 0 \\ [-1,-1,1,2]_1 &= 149/300 - 41/(150u) + 1/\hat{e}_1 * (-1/5 + 1/(10u)) \\ [-1,-2,2,2]_1 &= 1/5 - 77/(450u^2) + 26/(75u) + 1/\hat{e}_1 * (1/(15u^2) - 1/(5u)) \end{aligned}$$

TABLE 4(E) $K=-1, \alpha_1=2$

$$[K,M,N,L]_1 = \oint d^4q ((p-q)^2)^K (q^2)^M (q^+)^N (q^-)^L q_i$$

all terms are coefficients of p_i

$$[-1,2,-1,0]_1 = (-u + 5u^2 - 5u^3) * \log u + u - 11u^2/2 + 11u^3/2 + 1/\hat{e}_1 * (-u + 5u^2 - 5u^3)$$

$$[-1,1,0,0]_1 = 0$$

$$[-1,0,1,0]_1 = 0$$

$$[-1,-1,2,0]_1 = 7/12 - 1/(4\hat{e}_1)$$

$$[-1,2,-1,1]_1 = (-u/2 + 2u^2 - 7u^3/4) * \log u + 3u/4 - 3u^2 + 41u^3/16 + 1/\hat{e}_1 * (-u/2 + 2u^2 - 7u^3/4)$$

$$[-1,1,0,1]_1 = 0$$

$$[-1,0,1,1]_1 = 0$$

$$[-1,-1,2,1]_1 = 149/300 - 41/(150u) + 1/\hat{e}_1 * (-1/5 + 1/(10u))$$

$$[-1,2,-1,2]_1 = -u/3 * \log u + 7u^2/6 * \log u - 14u^3/15 * \log u + 11u/18 - 151u^2/72 + 367u^3/225 + 1/\hat{e}_1 * (-u/3 + 7u^2/6 - 14u^3/15)$$

$$[-1,1,0,2]_1 = 0$$

$$[-1,0,1,2]_1 = 0$$

$$[-1,-1,2,2]_1 = 157/360 + 23/(450u^2) - 169/(450u) + 1/\hat{e}_1 * (-1/6 - 1/(60u^2) + 2/(15u))$$

TABLE 5(A) $K = -2$, $\alpha_1 = -2$

$$[K, M, N, L]_2 = f \int d^4 q ((p-q)^2)^K (q^2)^M (q^+)^N (q^-)^L q_i q_j$$

terms in parentheses () are coefficients of $p_i p_j$
 terms in braces { } are coefficients of $p^2 \delta(i, j)$

$$[-2, 0, -2, 0]_2 = (- 1 + \log u) + (1/\hat{e}_0) + \{ - u/2 \}$$

$$[-2, -1, -1, 0]_2 = (- 1 + \log u/(u-1)^2 - 1/(u-1)) + (1/\hat{e}_0) \\ - \{ u/2/(u-1)*\log u \}$$

$$[-2, -2, 0, 0]_2 = - (2) + (1/\hat{e}_0) - \{1/2\}$$

$$[-2, 0, -2, 1]_2 = (- 2 + \log u) + (1/\hat{e}_0) + \{ - u/4 \}$$

$$[-2, -1, -1, 1]_2 = (- 3/2 + (2-u)/2/(u-1)^2 * \log u - 1/2/(u-1) + 1/\hat{e}_0) \\ - \{ u/4/(u-1)*\log u \}$$

$$[-2, -2, 0, 1]_2 = - (5/2) + (1/\hat{e}_0) - \{1/4\}$$

$$[-2, 0, -2, 2]_2 = (- 5/2 + \log u) + (1/\hat{e}_0) + \{ - u/6 \}$$

$$[-2, -1, -1, 2]_2 = (- 11/6 + (3-2u)/3/(u-1)^2 * \log u - 1/3/(u-1)) + (1/\hat{e}_0) \\ - \{ u/6/(u-1)*\log u \}$$

$$[-2, -2, 0, 2]_2 = - (17/6) + (1/\hat{e}_0) - \{1/6\}$$

TABLE 5(B) $K = -2$, $\alpha_1 = -1$

$$[K,M,N,L]_2 = f \int d^4q ((p-q)^2)^K (q^2)^M (q^+)^N (q^-)^L q_i q_j$$

terms in parentheses () are coefficients of $p_i p_j$

terms in braces { } are coefficients of $p^2 \delta(i,j)$

$$[-2,1,-2,0]_2 = (- 5/2 + \log u + 3u/2 - 3/2/(u-1)) + (1/\hat{e}_0)$$

$$+ \{ 3u^2/4 * \log u - u/2 + 13u^2/8 \} + \{ 1/\hat{e}_1 \} * \{ 3u^2/4 \}$$

$$[-2,0,-1,0]_2 = (- 1 + \log u - 1/(u-1)) + (1/\hat{e}_0) + \{ u/2 * \log u + u/2 \} + \{ 1/\hat{e}_1 \} * \{ u/2 \}$$

$$[-2,-1,0,0]_2 = (- 3/2 - 1/2/(u-1)) + (1/\hat{e}_0) + \{ 1/4 \} * \{ 1/\hat{e}_1 \}$$

$$[-2,-2,1,0]_2 = - (5/2) + (1/\hat{e}_0) - \{ 1/4 \}$$

$$[-2,1,-2,1]_2 = (- 8/3 + \log u + 4u/3 - 2/(3(u-1))) + (1/\hat{e}_0)$$

$$+ \{ u^2/3 * \log u - u/4 + 5u^2/9 \} + \{ 1/\hat{e}_1 \} * \{ u^2/3 \}$$

$$[-2,0,-1,1]_2 = (- 3/2 + \log u - 1/2/(u-1)) + (1/\hat{e}_0)$$

$$+ \{ u/4 * \log u + u/8 \} + \{ 1/\hat{e}_1 \} * \{ u/4 \}$$

$$[-2,-1,0,1]_2 = (- 11/6 - 1/(3(u-1))) + (1/\hat{e}_0) - \{ 1/36 \} + \{ 1/6 \} * \{ 1/\hat{e}_1 \}$$

$$[-2,-2,1,1]_2 = (- 17/6 + 1/2/u - 1/6/(u-1)) + (1/\hat{e}_0)$$

$$+ \{ - 1/6 + 1/36/u \} + \{ 1/\hat{e}_1 \} * \{ 1/12/u \}$$

$$[-2,1,-2,2]_2 = (- 35/12 + \log u + 5u/4 - 5/12/(u-1)) + (1/\hat{e}_0)$$

$$+ \{ 5u^2/24 * \log u - u/6 + 79u^2/288 \} + \{ 1/\hat{e}_1 \} * \{ 5u^2/24 \}$$

$$[-2,0,-1,2]_2 = (- 11/6 + \log u - 1/(3(u-1))) + (1/\hat{e}_0) \\ + \{ u/6 \log u + u/36 \} + \{ 1/\hat{e}_1 \} * \{ u/6 \}$$

$$[-2,-1,0,2]_2 = (- 25/12 - 1/4/(u-1)) + (1/\hat{e}_0) - \{ 1/24 \} + \{ 1/8 \} * \{ 1/\hat{e}_1 \}$$

$$[-2,-2,1,2]_2 = (- 37/12 + 2/3 u - 1/6/(u-1) + 1/\hat{e}_0) - \{ 1/8 - 1/36 u \} \\ + \{ 1/\hat{e}_1 \} * \{ 1/12 u \}$$

TABLE 5(C) $K = -2$, $\alpha_1 = 0$

$$[K, M, N, L]_2 = f \int d^4 q ((p-q)^2)^K (q^2)^M (q^+)^N (q^-)^L q_i q_j$$

terms in parentheses () are coefficients of $p_i p_j$
 terms in braces { } are coefficients of $p^2 \delta(i, j)$

$$\begin{aligned} [-2, 2, -2, 0]_2 &= (5/3 + (1-10u^2)*\log u + 32u/3 + 8u^2/3 + 8/3/(u-1) + 1/\hat{e}_0) \\ &+ (1/\hat{e}_1) * (-10u^2) + \{1/\hat{e}_1\} * \{2u^2 - 10u^3/3\} \\ &+ \{2u^2*\log u - 10u^3/3*\log u - u/2 + 4u^2 - 47u^3/9\} \end{aligned}$$

$$\begin{aligned} [-2, 1, -1, 0]_2 &= (1 + \log u - 4u*\log u + 4u + 1/(u-1)) + (1/\hat{e}_0) \\ &+ (1/\hat{e}_1) * (-4u) + \{1/\hat{e}_1\} * \{u/2 - u^2\} \\ &+ \{u/2*\log u - u^2*\log u + u/2 - u^2\} \end{aligned}$$

$$[-2, 0, 0, 0]_2 = (1/\hat{e}_0) - (1/\hat{e}_1)$$

$$[-2, -1, 1, 0]_2 = (-11/6 - 1/(3(u-1))) + (1/\hat{e}_0) - \{1/36\} + \{1/6\} * \{1/\hat{e}_1\}$$

$$[-2, -2, 2, 0]_2 = - (17/6) + (1/\hat{e}_0) - \{1/6\}$$

$$\begin{aligned} [-2, 2, -2, 1]_2 &= (-7/6 + (1-5u^2)*\log u + 35u/6 + 3u^2 + 5/6/(u-1) + 1/\hat{e}_0) \\ &+ (1/\hat{e}_1) * (-5u^2) + \{1/\hat{e}_1\} * \{5u^2/6 - 5u^3/4\} \\ &+ \{5u^2/6*\log u - 5u^3/4*\log u - u/4 + 47u^2/36 - 23u^3/16\} \end{aligned}$$

$$\begin{aligned}
[-2,1,-1,1]_2 &= (- 2/3 + \log u - 5u/2*\log u + 43u/12 + 1/3/(u-1) + 1/\hat{e}_0) \\
&+ (1/\hat{e}_1) * (- 5u/2) + \{1/\hat{e}_1\} * \{ u/4 - 5u^2/12 \} \\
&+ \{ u/4*\log u - 5u^2/12*\log u + u/8 - 17u^2/72 \}
\end{aligned}$$

$$[-2,0,0,1]_2 = (1/\hat{e}_0) - (1/\hat{e}_1)$$

$$\begin{aligned}
[-2,-1,1,1]_2 &= (- 25/12 + 1/2/u - 1/6/(u-1)) + (1/\hat{e}_0) + (1/\hat{e}_1) * (- 1/4/u) \\
&+ \{ - 1/24 + 1/36/u \} + \{1/\hat{e}_1\} * \{ 1/8 - 1/24/u \}
\end{aligned}$$

$$\begin{aligned}
[-2,-2,2,1]_2 &= (- 37/12 + 2/3/u - 1/6/(u-1)) + (1/\hat{e}_0) \\
&+ \{ - 1/8 + 1/36/u \} + \{1/\hat{e}_1\} * \{ 1/12/u \}
\end{aligned}$$

$$\begin{aligned}
[-2,2,-2,2]_2 &= (- 21/10 + \log u - 7u^2/2*\log u + 22u/5 + 121u^2/40 + 2/5/(u-1) + 1/\hat{e}_0) \\
&+ (1/\hat{e}_1) * (- 7u^2/2) + \{1/\hat{e}_1\} * \{ u^2/2 - 7u^3/10 \} \\
&+ \{ u^2/2*\log u - 7u^3/10*\log u - u/6 + 5u^2/8 - 117u^3/200 \}
\end{aligned}$$

$$\begin{aligned}
[-2,1,-1,2]_2 &= (- 4/3 + \log u - 2u*\log u + 7u/2 + 1/6/(u-1) + 1/\hat{e}_0) \\
&+ (1/\hat{e}_1) * (- 2u) + \{1/\hat{e}_1\} * \{ u/6 - u^2/4 \} \\
&+ \{ u/6*\log u - u^2/4*\log u + u/36 - u^2/16 \}
\end{aligned}$$

$$[-2,0,0,2]_2 = (1/\hat{e}_0) - (1/\hat{e}_1)$$

$$\begin{aligned}
[-2,-1,1,2]_2 &= (- 137/60 + 67/75/u - 1/10/(u-1)) + (1/\hat{e}_0) + (1/\hat{e}_1) * (- 2/5/u) \\
&+ \{ - 29/600 + 11/300/u \} + \{1/\hat{e}_1\} * \{ 1/10 - 1/20/u \}
\end{aligned}$$

$$\begin{aligned}
[-2,-2,2,2]_2 &= (- 197/60 + 8/75/u^2 + 14/15/u - 2/15/(u-1) + 1/\hat{e}_0) \\
&+ (1/\hat{e}_1) * (- 1/10/u^2) + \{1/\hat{e}_1\} * \{ - 1/30/u^2 + 1/10/u \} \\
&+ \{ - 1/10 + 17/900/u^2 + 2/75/u \}
\end{aligned}$$

TABLE 5(D) $K = -2$, $\alpha_1 = 0$

$$[K,M,N,L]_2 = f \int d^4q ((p-q)^2)^K (q^2)^M (q^+)^N (q^-)^L q_i q_j$$

terms in parentheses () are coefficients of $p_i p_j$
 terms in braces {} are coefficients of $p^2 \delta(i,j)$

$$\begin{aligned} [-2,2,-1,0]_2 &= (- u/(u-1) + (1-10u+15u^2)*\log u + 6u - 33u^2/2 + 1/\hat{e}_0) \\ &+ (1/\hat{e}_1) * (- 10u + 15u^2) + \{1/\hat{e}_1\} * \{ u/2 - 5u^2/2 + 5u^3/2 \} \\ &+ \{ u/2*\log u + 5u^2(u-1)/2*\log u + u/2 - 9u^2/4 + 9u^3/4 \} \end{aligned}$$

$$[-2,1,0,0]_2 = (1/\hat{e}_0) - (1/\hat{e}_1)$$

$$[-2,0,1,0]_2 = (1/\hat{e}_0) - (1/\hat{e}_1)$$

$$[-2,-1,2,0]_2 = (- 25/12 - 1/4/(u-1)) + (1/\hat{e}_0) - \{1/24\} + \{1/8\} * \{1/\hat{e}_1\}$$

$$\begin{aligned} [-2,2,-1,1]_2 &= (- 5/4 + (1-6u+7u^2)*\log u + 29u/4 - 41u^2/4 - 1/4/(u-1)) + (1/\hat{e}_0) \\ &+ (1/\hat{e}_1) * (- 6u + 7u^2) + \{1/\hat{e}_1\} * \{ u/4 - u^2 + 7u^3/8 \} \\ &+ \{ (u/4 - u^2 + 7u^3/8)*\log u + u/8 - u^2/2 + 15u^3/32 \} \end{aligned}$$

$$[-2,1,0,1]_2 = (1/\hat{e}_0) - (1/\hat{e}_1)$$

$$[-2,0,1,1]_2 = (1/\hat{e}_0) - (1/\hat{e}_1)$$

$$\begin{aligned} [-2,-1,2,1]_2 &= (- 137/60 + 67/75/u - 1/10/(u-1)) + (1/\hat{e}_0) + (1/\hat{e}_1) * (- 2/5/u) \\ &+ \{ - 29/600 + 11/300/u \} + \{1/\hat{e}_1\} * \{ 1/10 - 1/20/u \} \end{aligned}$$

$$\begin{aligned}
[-2,2,-1,2]_2 &= (- 8/5 + (1-14u/3+14u^2/3)*\log u + 671u/90 - 367u^2/45 - 1/10/(u-1)) + (1/\hat{e}_0) \\
&+ (1/\hat{e}_1) * (- 14u/3 + 14u^2/3) + \{1/\hat{e}_1\} * \{ u/6 - 7u^2/12 + 7u^3/15 \} \\
&+ \{ (u/6-7u^2/12+7u^3/15)*\log u + u/36 - 17u^2/144 + 53u^3/450 \} \\
\\
[-2,1,0,2]_2 &= (1/\hat{e}_0) - (1/\hat{e}_1) \\
[-2,0,1,2]_2 &= (1/\hat{e}_0) - (1/\hat{e}_1) \\
[-2,-1,2,2]_2 &= (- 49/20 - 77/450/u^2 + 293/180/u - 1/20/(u-1)) + (1/\hat{e}_0) \\
&+ (1/\hat{e}_1) * (1/15/u^2 - 2/3/u) + \{1/\hat{e}_1\} * \{ 1/12 + 1/120/u^2 - 1/15/u \} \\
&+ \{ - 37/720 - 2/225/u^2 + 49/900/u \}
\end{aligned}$$

TABLE 5(E) $K = -2$, $\alpha_1 = 2$

This Table is identical to Table 1(E) with the understanding that all terms are enclosed in parentheses ().

TABLE 6(A) $K = -1$, $\alpha_1 = -2$

$$[K,M,N,L]_2 = \int d^4q ((p-q)^2)^K (q^2)^M (q^+)^N (q^-)^L q_i q_j$$

terms in parentheses () are coefficients of $p_i p_j$

terms in braces {} are coefficients of $p^2 \delta(i,j)$

$$[-1,-1,-2,0]_2 = (-1 + \log u + \log u/(u-1)^2 + 2/(u-1)*\log u - 1/(u-1)) \\ + \{ -1/2*\log u - 1/2/(u-1)*\log u - u/2 - S_1/2 \}$$

$$[-1,-2,-1,0]_2 = (1 - \log u/(u-1)^2 - \log u/(u-1) + 1/(u-1)) \\ + \{ 1/2*\log u + 1/2/(u-1)*\log u + S_1/2 \}$$

$$[-1,-1,-2,1]_2 = (-u/2/(u-1) + u^2/2/(u-1)^2*\log u) \\ + \{ -1/2 + 1/4*\log u - 1/4/(u-1)*\log u - u/8 - S_1/2/u \}$$

$$[-1,-2,-1,1]_2 = (1/2 - 1/2/(u-1)^2*\log u - 1/2/(u-1)*\log u + 1/2/(u-1)) \\ + \{ 1/2 - 1/4*\log u + 1/4/(u-1)*\log u + S_1/2/u \}$$

$$[-1,-1,-2,2]_2 = (-u/3/(u-1) + u^2/3/(u-1)^2*\log u) \\ - \{ 1/8 - (u^2+3u-6)/12/u/(u-1)*\log u + u/18 + S_1/2/u^2 + 1/2/u \}$$

$$[-1,-2,-1,2]_2 = (-u/3/(u-1)^2*\log u + u/3/(u-1)) \\ + \{ 1/8 - (u^2+3u-6)/12/u/(u-1)*\log u + S_1/2/u^2 + 1/2/u \}$$

TABLE 6(B) K = -1 , $\alpha_1 = -1$

$$[K,M,N,L]_2 = f \int d^4q ((p-q)^2)^K (q^2)^M (q^+)^N (q^-)^L q_i q_j$$

terms in parentheses () are coefficients of $p_i p_j$
 terms in braces { } are coefficients of $p^2 \delta(i,j)$

$$[-1,0,-2,0]_2 = (- 1/2 + u/2 - 1/2/(u-1)) \\ + \{ u^2/4 * \log u + 3u^2/8 \} + \{ 1/\hat{e}_1 \} * \{ u^2/4 \}$$

$$[-1,-1,-1,0]_2 = (1/2 * \log u - 1/2/(u-1)^2 * \log u) \\ + \{ u^2/4/(u-1) * \log u \} + \{ 1/\hat{e}_1 \} * \{ u/4 \}$$

$$[-1,-2,0,0]_2 = (1/2 - 1/2/(u-1)) + \{ 1/4 \} * \{ 1/\hat{e}_1 \}$$

$$[-1,0,-2,1]_2 = (- 1/6 + u/3 - 1/6/(u-1)) \\ + \{ u^2/12 * \log u + 7u^2/72 \} + \{ 1/\hat{e}_1 \} * \{ u^2/12 \}$$

$$[-1,-1,-1,1]_2 = (1/3 * \log u - 1/6/(u-1)^2 * \log u + 1/6/(u-1) * \log u) \\ + \{ u^2/12/(u-1) * \log u \} - u/72 \} + \{ 1/\hat{e}_1 \} * \{ u/12 \}$$

$$[-1,-2,0,1]_2 = (1/3 - 1/6/(u-1)) + \{ 1/36 \} + \{ 1/12 \} * \{ 1/\hat{e}_1 \}$$

$$[-1,0,-2,2]_2 = (- 1/12 + u/4 - 1/12/(u-1)) \\ + \{ u^2/24 * \log u + 11u^2/288 \} + \{ 1/\hat{e}_1 \} * \{ u^2/24 \}$$

$$[-1,-1,-1,2]_2 = (1/4 * \log u - 1/12/(u-1)^2 * \log u + 1/6/(u-1) * \log u) \\ + \{ u^2/24/(u-1) * \log u - u/72 \} + \{ 1/\hat{e}_1 \} * \{ u/24 \}$$

$$[-1,-2,0,2]_2 = (1/4 - 1/12/(u-1)) + \{ 1/72 \} + \{ 1/24 \} * \{ 1/\hat{e}_1 \}$$

TABLE 6(C) $K = -1$, $\alpha_1 = 0$

$$[K, M, N, L]_2 = f \int d^4 q ((p-q)^2)^K (q^2)^M (q^+)^N (q^-)^L q_i q_j$$

terms in parentheses () are coefficients of $p_i p_j$
 terms in braces {} are coefficients of $p^2 \delta(i, j)$

$$\begin{aligned} [-1, 1, -2, 0]_2 &= (5/6 - 2u^2 \log u + 11u/6 + 4u^2/3 + 5/6/(u-1)) \\ &+ (1/\hat{e}_1) * (- 2u^2) + \{1/\hat{e}_1\} * \{ u^2/4 - 2u^3/3 \} \\ &+ \{ u^2/4 \log u - 2u^3/3 \log u + 3u^2/8 - 7u^3/9 \} \\ [-1, 0, -1, 0]_2 &= (1/2 - u \log u + 3u/2 + 1/2/(u-1)) + (1/\hat{e}_1) * (- u) \\ &+ \{ -u^2/4 \log u - u^2/8 \} + \{1/\hat{e}_1\} * \{ - u^2/4 \} \\ [-1, -1, 0, 0]_2 &= (13/18 + 1/6/(u-1) - 1/3/\hat{e}_1) + \{1/18\} - \{1/12\} * \{1/\hat{e}_1\} \\ [-1, -2, 1, 0]_2 &= (1/3 - 1/6/(u-1)) + \{1/36\} + \{1/12\} * \{1/\hat{e}_1\} \\ [-1, 1, -2, 1]_2 &= (1/4 - 5u^2/6 \log u + 3u/4 + 7u^2/9 + 1/4/(u-1)) \\ &+ (1/\hat{e}_1) * (- 5u^2/6) + \{1/\hat{e}_1\} * \{ u^2/12 - 5u^3/24 \} \\ &+ \{ u^2/12 \log u - 5u^3/24 \log u + 7u^2/72 - 49u^3/288 \} \\ [-1, 0, -1, 1]_2 &= (- u/2 \log u + 11u/12 + u/6/(u-1)) - (1/\hat{e}_1) * (u/2) \\ &+ \{ - u^2/12 \log u - u^2/72 \} + \{1/\hat{e}_1\} * \{ - u^2/12 \} \\ [-1, -1, 0, 1]_2 &= (7/12 + 1/12/(u-1) - 1/4/\hat{e}_1) + \{1/36\} - \{1/24\} * \{1/\hat{e}_1\} \\ [-1, -2, 1, 1]_2 &= (1/4 + 1/18/u) + (1/\hat{e}_1) * (- 1/12/u) \\ &+ \{ 1/72 + 1/36/u \} + \{1/\hat{e}_1\} * \{ 1/24 - 1/24/u \} \end{aligned}$$

$$\begin{aligned}
[-1,1,-2,2]_2 &= (7/60 - u^2/2 \cdot \log u + 9u/20 + 23u^2/40 + 7/60/(u-1)) \\
&+ (1/\hat{e}_1) * (- u^2/2) + \{1/\hat{e}_1\} * \{ u^2/24 - u^3/10 \} \\
&+ \{ u^2/24 \cdot \log u - u^3/10 \cdot \log u + 11u^2/288 - 11u^3/200 \}
\end{aligned}$$

$$\begin{aligned}
[-1,0,-1,2]_2 &= (-u/3 \cdot \log u + 25u/36 + u/12/(u-1)) - (1/\hat{e}_1) * (u/3) \\
&+ \{ - u^2/24 \cdot \log u + u^2/288 \} + \{1/\hat{e}_1\} * \{ - u^2/24 \}
\end{aligned}$$

$$[-1,-1,0,2]_2 = (149/300 + 1/20/(u-1)) - (1/5/\hat{e}_1) + \{11/600\} - \{1/40/\hat{e}_1\}$$

$$\begin{aligned}
[-1,-2,1,2]_2 &= (1/5 + 8/75/u + 1/60/(u-1)) + (1/\hat{e}_1) * (- 1/10/u) \\
&+ \{ 1/150 + 17/900/u \} + \{1/\hat{e}_1\} * \{ 1/40 - 1/30/u \}
\end{aligned}$$

TABLE 6(D) $K = -1$, $\alpha_1 = 1$

$$[K,M,N,L]_2 = f \int d^4q ((p-q)^2)^K (q^2)^M (q^+)^N (q^-)^L q_i q_j$$

terms in parentheses () are coefficients of $p_i p_j$

terms in braces {} are coefficients of $p^2 \delta(i,j)$

$$\begin{aligned} [-1,2,-2,0]_2 &= (-11u/12/(u-1) - (5u^2-10u^3)*\log u + u/12 + u^2/12 - 39u^3/4) \\ &+ (1/\hat{e}_1) * (-5u^2 + 10u^3) + \{1/\hat{e}_1\} * \{ u^2/4 - 5u^3/3 + 15u^4/8 \} \\ &+ \{ (u^2/4-5u^3/3+15u^4/8)*\log u + 3u^2/8 - 16u^3/9 + 59u^4/32 \} \end{aligned}$$

$$\begin{aligned} [-1,1,-1,0]_2 &= (-u/3/(u-1) - u*\log u + 5u^2/2*\log u + 2u/3 - 43u^2/12) \\ &+ (1/\hat{e}_1) * (-u + 5u^2/2) + \{1/\hat{e}_1\} * \{ -u^2/4 + 5u^3/12 \} \\ &+ \{ -u^2/4*\log u + 5u^3/12*\log u - u^2/8 + 17u^3/72 \} \end{aligned}$$

$$[-1,0,0,0]_2 = 0$$

$$[-1,-1,1,0]_2 = (7/12 + 1/12/(u-1) - 1/4/\hat{e}_1) + \{1/36\} - \{1/24\} * \{1/\hat{e}_1\}$$

$$[-1,-2,2,0]_2 = (1/4 - 1/12/(u-1)) + \{1/72\} + \{1/24\} * \{1/\hat{e}_1\}$$

$$\begin{aligned} [-1,2,-2,1]_2 &= (-13u/60/(u-1) - (2u^2-7u^3/2)*\log u + 17u/60 + 19u^2/20 - 177u^3/40) \\ &+ (1/\hat{e}_1) * (-2u^2 + 7u^3/2) + \{1/\hat{e}_1\} * \{ u^2/12 - u^3/2 + 21u^4/40 \} \\ &+ \{ (u^2/12-u^3/2+21u^4/40)*\log u + 7u^2/72 - 3u^3/8 + 281u^4/800 \} \end{aligned}$$

$$\begin{aligned} [-1,1,-1,1]_2 &= (-u/12/(u-1) - u/2*\log u + u^2*\log u + 2u/3 - 7u^2/4) \\ &+ (1/\hat{e}_1) * (-u/2 + u^2) + \{1/\hat{e}_1\} * \{ -u^2/12 + u^3/8 \} \\ &+ \{ -u^2/12*\log u + u^3/8*\log u - u^2/72 + u^3/32 \} \end{aligned}$$

$$[-1,0,0,1]_2 = 0$$

$$\begin{aligned} [-1,-1,1,1]_2 &= (149/300 - 3/25/u + 1/30/(u-1)) - (1/\hat{e}_1) * (1/5 - 1/20/u) \\ &+ \{ 11/600 - 2/225/u \} + \{1/\hat{e}_1\} * \{ - 1/40 + 1/120/u \} \end{aligned}$$

$$\begin{aligned} [-1,-2,2,1]_2 &= (1/5 + 8/75/u + 1/60/(u-1)) + (1/\hat{e}_1) * (- 1/10/u) \\ &+ \{ 1/150 + 17/900/u \} + \{1/\hat{e}_1\} * \{ 1/40 - 1/30/u \} \end{aligned}$$

$$\begin{aligned} [-1,2,-2,2]_2 &= (-u/12/(u-1) - (7u^2/6 - 28u^3/15)*\log u + u/4 + 67u^2/72 - 629u^3/225) \\ &+ (1/\hat{e}_1) * (- 7u^2/6 + 28u^3/15) \\ &+ \{ (u^2/24 - 7u^3/30 + 7u^4/30)*\log u + 11u^2/288 - 211u^3/1800 + 22u^4/225 \} \\ &+ \{1/\hat{e}_1\} * \{ u^2/24 - 7u^3/30 + 7u^4/30 \} \end{aligned}$$

$$\begin{aligned} [-1,1,-1,2]_2 &= (- u/30/(u-1) - (u/3 - 7u^2/12)*\log u + 26u/45 - 839u^2/720) \\ &+ (1/\hat{e}_1) * (- u/3 + 7u^2/12) + \{1/\hat{e}_1\} * \{ - u^2/24 + 7u^3/120 \} \\ &+ \{ - u^2/24*\log u + 7u^3/120*\log u + u^2/288 + u^3/7200 \} \end{aligned}$$

$$[-1,0,0,2]_2 = 0$$

$$\begin{aligned} [-1,-1,1,2]_2 &= (157/360 - 77/450/u + 1/60/(u-1)) \\ &+ (1/\hat{e}_1) * (- 1/6 + 1/15/u) + \{ 49/3600 - 2/225/u \} \\ &+ \{1/\hat{e}_1\} * \{ - 1/60 + 1/120/u \} \end{aligned}$$

$$\begin{aligned} [-1,-2,2,2]_2 &= (1/6 - 31/450/u^2 + 173/900/u + 1/60/(u-1)) \\ &+ (1/\hat{e}_1) * (1/30/u^2 - 2/15/u) + \{1/\hat{e}_1\} * \{ 1/60 + 1/120/u^2 - 1/30/u \} \\ &+ \{ 11/3600 - 2/225/u^2 + 17/900/u \} \end{aligned}$$

TABLE 6(E) $K = -1$, $\alpha_1 = 2$

$$[K,M,N,L]_2 = f \int d^4q ((p-q)^2)^K (q^2)^M (q^+)^N (q^-)^L q_i q_j$$

terms in parentheses () are coefficients of $p_i p_j$

terms in braces { } are coefficients of $p^2 \delta(i,j)$

$$\begin{aligned} [-1,2,-1,0]_2 &= (u/4/(u-1) - (u-6u^2+7u^3)*\log u + 5u/4 - 29u^2/4 + 41u^3/4) \\ &+ (1/\hat{e}_1) * (-u + 6u^2 - 7u^3) + \{1/\hat{e}_1\} * \{ -u^2/4 + u^3 - 7u^4/8 \} \\ &+ \{ (-u^2/4 + u^3 - 7u^4/8)*\log u - u^2/8 + u^3/2 - 15u^4/32 \} \end{aligned}$$

$$[-1,1,0,0]_2 = 0$$

$$[-1,0,1,0]_2 = 0$$

$$[-1,-1,2,0]_2 = (149/300 + 1/20/(u-1)) - (1/5/\hat{e}_1) + \{11/600\} - \{1/40/\hat{e}_1\}$$

$$\begin{aligned} [-1,2,-1,1]_2 &= (- (u/2-7u^2/3+7u^3/3)*\log u + 4u/5 - 671u^2/180 + 367u^3/90 + u/20/(u-1)) \\ &+ (1/\hat{e}_1) * (-u/2 + 7u^2/3 - 7u^3/3) - \{1/\hat{e}_1\} * \{ u^2/12 - 7u^3/24 + 7u^4/30 \} \\ &+ \{ -(u^2/12-7u^3/24+7u^4/30)*\log u - u^2/72 + 17u^3/288 - 53u^4/900 \} \end{aligned}$$

$$[-1,1,0,1]_2 = 0$$

$$[-1,0,1,1]_2 = 0$$

$$\begin{aligned} [-1,-1,2,1]_2 &= (157/360 - 77/450/u + 1/60/(u-1)) + (1/\hat{e}_1) * (-1/6 + 1/15/u) \\ &+ \{ 49/3600 - 2/225/u \} + \{1/\hat{e}_1\} * \{ -1/60 + 1/120/u \} \end{aligned}$$

$$\begin{aligned}
[-1,2,-1,2]_2 &= (-(u/3-4u^2/3+6u^3/5)*\log u +113u/180 -113u^2/45 +178u^3/75 +u/60/(u-1)) \\
&+ (1/\hat{e}_1) * (-u/3 + 4u^2/3 - 6u^3/5) - \{1/\hat{e}_1\} * \{ u^2/24 - 2u^3/15 + u^4/10 \} \\
&+ \{ -(u^2/24-2u^3/15+u^4/10)*\log u +u^2/288 - u^3/225 - u^4/450\}
\end{aligned}$$

$$[-1,1,0,2]_2 = 0$$

$$[-1,0,1,2]_2 = 0$$

$$\begin{aligned}
[-1,-1,2,2]_2 &= (383/980 + 599/22050/u^2 - 2291/8820/u + 1/140/(u-1)) \\
&- (1/\hat{e}_1) * (1/7 +1/105/u^2 -2/21/u) - \{1/\hat{e}_1\} * \{ 1/84 +1/840/u^2 - 1/105/u \} \\
&+ \{ 379/35280 + 71/44100/u^2 - 463/44100/u \}
\end{aligned}$$

TABLE 7(A) $K = -2$, $\alpha_1 = -2$

$$[K,M,N,L]_3 = f \int d^4 q ((p-q)^2)^K (q^2)^M (q^+)^N (q^-)^L q_i q_j q_k$$

terms in parentheses () are coefficients of $p_i p_j p_k$

terms in braces { } are coefficients of $p_i^2 \delta(j,k)$ plus two cyclic permutations

$$[-2,0,-2,0]_3 = (- 1 + \log u) + (1/\hat{e}_0) + \{ - u/2 \}$$

$$[-2,-1,-1,0]_3 = (- 3/2 - 1/(u-1)^3 \log u + 1/(u-1)^2 - 1/2/(u-1)) \\ + (1/\hat{e}_0) + \{ - 1/4 - 1/4 \log u + 1/4/(u-1)^2 \log u - 1/4/(u-1) \}$$

$$[-2,-2,0,0]_3 = - (5/2) + (1/\hat{e}_0) - \{1/4\}$$

$$[-2,0,-2,1]_3 = (- 2 + \log u) + (1/\hat{e}_0) + \{ - u/4 \}$$

$$[-2,-1,-1,1]_3 = (- 11/6 - 1/3/(u-1)^3 \log u + 1/3/(u-1)^2 \log u - 1/3/(u-1) \log u + 1/3/(u-1)^2 \\ - 1/2/(u-1)) + (1/\hat{e}_0) \\ + \{ - 1/12 - 1/6 \log u + 1/12/(u-1)^2 \log u - 1/12/(u-1) \log u - 1/12/(u-1) \}$$

$$[-2,-2,0,1]_3 = - (17/6) + (1/\hat{e}_0) - \{1/6\}$$

$$[-2,0,-2,2]_3 = (- 5/2 + \log u) + (1/\hat{e}_0) + \{ - u/6 \}$$

$$[-2,-1,-1,2]_3 = (- 25/12 - 1/6/(u-1)^3 \log u + 1/3/(u-1)^2 \log u - 1/2/(u-1) \log u + 1/6/(u-1)^2 \\ - 5/12/(u-1)) + (1/\hat{e}_0) \\ + \{ - 1/24 - 1/8 \log u + 1/24/(u-1)^2 \log u - 1/12/(u-1) \log u - 1/24/(u-1) \}$$

$$[-2,-2,0,2]_3 = - (37/12) + (1/\hat{e}_0) - \{1/8\}$$

TABLE 7(B) $K = -2$, $\alpha_1 = -1$

$$[K,M,N,L]_3 = f \int d^4q ((p-q)^2)^K (q^2)^M (q^+)^N (q^-)^L q_i q_j q_k$$

terms in parentheses () are coefficients of $p_i p_j p_k$

terms in braces { } are coefficients of $p_i^2 \delta(j,k)$ plus two cyclic permutations

$$[-2,1,-2,0]_3 = (- 7 + \log u - 2u - 6/(u-1)) + (1/\hat{e}_0) \\ + \{ u^2 * \log u - u/2 + 2u^2 \} + \{ 1/\hat{e}_1 * (u^2) \}$$

$$[-2,0,-1,0]_3 = (- 3 + \log u - 3/(u-1)) + (1/\hat{e}_0) \\ + \{ u/2 * \log u + u/2 \} + \{ 1/\hat{e}_1 * (u/2) \}$$

$$[-2,-1,0,0]_3 = (- 11/6 - 1/(u-1)) + (1/\hat{e}_0) - \{1/36\} + 1/6 * \{ 1/\hat{e}_1 \}$$

$$[-2,-2,1,0]_3 = - (17/6) + (1/\hat{e}_0) - \{1/6\}$$

$$[-2,1,-2,1]_3 = (- 9/2 + \log u - 5/2/(u-1)) + (1/\hat{e}_0) \\ + \{ 5u^2/12 * \log u - u/4 + 47u^2/72 \} + \{ 1/\hat{e}_1 * (5u^2/12) \}$$

$$[-2,0,-1,1]_3 = (- 5/2 + \log u - 3/2/(u-1)) + (1/\hat{e}_0) + \{ u/4 * \log u + u/8 \} + \{ 1/\hat{e}_1 * (u/4) \}$$

$$[-2,-1,0,1]_3 = (- 25/12 - 3/4/(u-1)) + (1/\hat{e}_0) - \{1/24\} + 1/8 * \{ 1/\hat{e}_1 \}$$

$$[-2,-2,1,1]_3 = (- 37/12 + 1/2/u - 1/4/(u-1)) + (1/\hat{e}_0) + \{ - 1/8 + 1/72/u \} + \{ 1/\hat{e}_1 * (1/24/u) \}$$

$$[-2,1,-2,2]_3 = (- 4 + \log u + u/2 - 3/2/(u-1)) + (1/\hat{e}_0) \\ + \{ u^2/4 * \log u - u/6 + 5u^2/16 \} + \{ 1/\hat{e}_1 * (u^2/4) \}$$

$$[-2,0,-1,2]_3 = (- 5/2 + \log u - 1/(u-1)) + (1/\hat{e}_0) + \{ u/6 * \log u + u/36 \} + \{ 1/\hat{e}_1 * (u/6) \}$$

$$[-2,-1,0,2]_3 = (- 137/60 - 3/5/(u-1)) + (1/\hat{e}_0) - \{29/600\} + 1/10 * \{ 1/\hat{e}_1 \}$$

$$[-2,-2,1,2]_3 = (- 197/60 + 7/10/u - 3/10/(u-1)) + (1/\hat{e}_0) \\ + \{ - 1/10 + 1/75/u \} + \{ 1/\hat{e}_1 * (1/20/u) \}$$

TABLE 7(C) $K = -2$, $\alpha_1 = 0$

$$[K,M,N,L]_3 = \int d^4q ((p-q)^2)^K (q^2)^M (q^+)^N (q^-)^L q_i q_j q_k$$

terms in parentheses () are coefficients of $p_i p_j p_k$

terms in braces { } are coefficients of $p_i^2 \delta(j,k)$ plus two cyclic permutations

$$\begin{aligned} [-2,2,-2,0]_3 &= (14 + \log u - 15u^2 \log u + 25u + 53u^2/2 + 15/(u-1)) \\ &+ (1/\hat{e}_0) + (1/\hat{e}_1 * (- 15u^2)) + \{ 1/\hat{e}_1 * (5u^2/2 - 5u^3) \} \\ &+ \{ 5u^2/2 \log u - 5u^3 \log u - u/2 + 19u^2/4 - 7u^3 \} \end{aligned}$$

$$\begin{aligned} [-2,1,-1,0]_3 &= (9/2 + \log u - 5u \log u + 2u/2 + 9/2/(u-1)) \\ &+ (1/\hat{e}_0) + (1/\hat{e}_1 * (- 5u)) + \{ 1/\hat{e}_1 * (u/2 - 5u^2/4) \} \\ &+ \{ u/2 \log u - 5u^2/4 \log u + u/2 - 9u^2/8 \} \end{aligned}$$

$$[-2,0,0,0]_3 = (1/\hat{e}_0) - (1/\hat{e}_1)$$

$$[-2,-1,1,0]_3 = (- 25/12 - 3/4/(u-1)) + (1/\hat{e}_0) - \{1/24\} + 1/8 * \{ 1/\hat{e}_1 \}$$

$$[-2,-2,2,0]_3 = - (37/12) + (1/\hat{e}_0) - \{1/8\}$$

$$\begin{aligned} [-2,2,-2,1]_3 &= (5/2 + \log u - 7u^2 \log u + 2u/2 + 12u^2 + 9/2/(u-1)) \\ &+ (1/\hat{e}_0) + (1/\hat{e}_1 * (- 7u^2)) + \{ 1/\hat{e}_1 * (u^2 - 7u^3/4) \} \\ &+ \{ u^2 \log u - 7u^3/4 \log u - u/4 + 3u^2/2 - 29u^3/16 \} \end{aligned}$$

$$\begin{aligned} [-2,1,-1,1]_3 &= (1/2 + \log u - 3u \log u + 13u/2 + 3/2/(u-1)) + (1/\hat{e}_0) + (1/\hat{e}_1 * (- 3u)) \\ &+ \{ u/4 \log u - u^2/2 \log u + u/8 - u^2/4 \} + \{ 1/\hat{e}_1 * (u/4 - u^2/2) \} \end{aligned}$$

$$[-2,0,0,1]_3 = (1/\hat{e}_0) - (1/\hat{e}_1)$$

$$\begin{aligned} [-2,-1,1,1]_3 &= (- 137/60 + 26/75/u - 9/20/(u-1)) + (1/\hat{e}_0) + (1/\hat{e}_1 * (- 1/5/u)) \\ &+ \{ - 29/600 + 11/600/u \} + \{ 1/\hat{e}_1 * (1/10 - 1/40/u) \} \end{aligned}$$

$$\begin{aligned} [-2,-2,2,1]_3 &= (- 197/60 + 7/10/u - 3/10/(u-1)) + (1/\hat{e}_0) \\ &+ \{ - 1/10 + 1/75/u \} + \{ 1/\hat{e}_1 * (1/20/u) \} \end{aligned}$$

$$\begin{aligned}
[-2,2,-2,2]_3 &= (- 2/5 + \log u - 14u^2/3 \cdot \log u + 203u/30 + 367u^2/45 + 21/10/(u-1)) \\
&+ (1/\hat{e}_0) + (1/\hat{e}_1 * (- 14u^2/3)) + \{ 1/\hat{e}_1 * (7u^2/12 - 14u^3/15) \} \\
&+ \{ 7u^2/12 \cdot \log u - 14u^3/15 \cdot \log u - u/6 + 101u^2/144 - 158u^3/225 \}
\end{aligned}$$

$$\begin{aligned}
[-2,1,-1,2]_3 &= (- 3/4 + \log u - 7u/3 \cdot \log u + 193u/36 + 3/4/(u-1)) \\
&+ (1/\hat{e}_0) + (1/\hat{e}_1 * (- 7u/3)) + \{ 1/\hat{e}_1 * (u/6 - 7u^2/24) \} \\
&+ \{ u/6 \cdot \log u - 7u^2/24 \cdot \log u + u/36 - 17u^2/288 \}
\end{aligned}$$

$$[-2,0,0,2]_3 = (1/\hat{e}_0) - (1/\hat{e}_1)$$

$$\begin{aligned}
[-2,-1,1,2]_3 &= (- 49/20 + 121/180/u - 3/10/(u-1)) + (1/\hat{e}_0) + (1/\hat{e}_1 * (- 1/3/u)) \\
&+ \{ - 37/720 + 49/1800/u \} + \{ 1/\hat{e}_1 * (1/12 - 1/30/u) \}
\end{aligned}$$

$$\begin{aligned}
[-2,-2,2,2]_3 &= (- 69/20 + 19/900/u^2 + 29/30/u - 3/10/(u-1)) + (1/\hat{e}_0) + (1/\hat{e}_1 * (- 1/15/u^2)) \\
&+ \{ - 1/12 + 17/1800/u^2 + 11/900/u \} + \{ 1/\hat{e}_1 * (- 1/60/u^2 + 1/15/u) \}
\end{aligned}$$

TABLE 7(D) $K = -2$, $\alpha_1 = 1$

$$[K,M,N,L]_3 = f \int d^4q ((p-q)^2)^K (q^2)^M (q^+)^N (q^-)^L q_i q_j q_k$$

terms in parentheses () are coefficients of $p_i p_j p_k$

terms in braces {} are coefficients of $p_i^2 \delta(j,k)$ plus two cyclic permutations

$$\begin{aligned} [-2,2,-1,0]_3 &= (- 6 + \log u - 12u \log u + 21u^2 \log u + 5u - 79u^2/2 - 6/(u-1)) \\ &+ (1/\hat{e}_0) + (1/\hat{e}_1 * (- 12u + 21u^2)) + \{ 1/\hat{e}_1 * (u/2 - 3u^2 + 7u^3/2) \} \\ &+ \{ u/2 \log u - 3u^2 \log u + 7u^3/2 \log u + u/2 - 5u^2/2 + 11u^3/4 \} \end{aligned}$$

$$[-2,1,0,0]_3 = (1/\hat{e}_0) - (1/\hat{e}_1)$$

$$[-2,0,1,0]_3 = (1/\hat{e}_0) - (1/\hat{e}_1)$$

$$[-2,-1,2,0]_3 = (- 137/60 - 3/5/(u-1)) + (1/\hat{e}_0) - \{29/600\} + 1/10 * \{ 1/\hat{e}_1 \}$$

$$\begin{aligned} [-2,2,-1,1]_3 &= (- 5/2 + \log u - 7u \log u + 28u^2/3 \log u + 17u/2 - 172u^2/9 - 3/2/(u-1)) \\ &+ (1/\hat{e}_0) + (1/\hat{e}_1 * (- 7u + 28u^2/3)) + \{ 1/\hat{e}_1 * (u/4 - 7u^2/6 + 7u^3/6) \} \\ &+ \{ u/4 \log u - 7u^2/6 \log u + 7u^3/6 \log u + u/8 - 19u^2/36 + 19u^3/36 \} \end{aligned}$$

$$[-2,1,0,1]_3 = (1/\hat{e}_0) - (1/\hat{e}_1)$$

$$[-2,0,1,1]_3 = (1/\hat{e}_0) - (1/\hat{e}_1)$$

$$\begin{aligned} [-2,-1,2,1]_3 &= (- 49/20 + 121/180/u - 3/10/(u-1)) + (1/\hat{e}_0) + (1/\hat{e}_1 * (- 1/3/u)) \\ &+ \{ - 37/720 + 49/1800/u \} + \{ 1/\hat{e}_1 * (1/12 - 1/30/u) \} \end{aligned}$$

$$\begin{aligned} [-2,2,-1,2]_3 &= (- 21/10 + \log u - 16u/3 \log u + 6u^2 \log u + 398u/45 - 199u^2/15 - 3/5/(u-1)) \\ &+ (1/\hat{e}_0) + (1/\hat{e}_1 * (- 16u/3 + 6u^2)) + \{ 1/\hat{e}_1 * (u/6 - 2u^2/3 + 3u^3/5) \} \\ &+ \{ u/6 \log u - 2u^2/3 \log u + 3u^3/5 \log u + u/36 - u^2/9 + 17u^3/150 \} \end{aligned}$$

$$[-2,1,0,2]_3 = (1/\hat{e}_0) - (1/\hat{e}_1)$$

$$[-2,0,1,2]_3 = (1/\hat{e}_0) - (1/\hat{e}_1)$$

$$\begin{aligned}
 [-2,-1,2,2]_3 &= (- 363/140 - 967/8820/u^2 + 64/49/u - 6/35/(u-1)) \\
 &+ (1/\hat{e}_0) + (1/\hat{e}_1 * (1/21/u^2 - 4/7/u)) + \{ 1/\hat{e}_1 * (1/14 + 1/210/u^2 - 1/21/u) \} \\
 &+ \{ - 103/1960 - 463/88200/u^2 + 379/8820/u \}
 \end{aligned}$$

TABLE 7(E) $K = -2$, $\alpha_1 = 2$

$$[K,M,N,L]_3 = f \int d^4q ((p-q)^2)^K (q^2)^M (q^+)^N (q^-)^L q_i q_j q_k$$

terms in parentheses () are coefficients of $p_i p_j p_k$

terms in braces { } are coefficients of $p_i^2 p_j \delta(j,k)$ plus two cyclic permutations

This Table is identical to Table 1(E) with the understanding that all terms are enclosed in parentheses ().

TABLE 8(A) $K = -1$, $\alpha_1 = -2$

$$[K,M,N,L]_3 = f \int d^4q ((p-q)^2)^K (q^2)^M (q^+)^N (q^-)^L q_i q_j q_k$$

terms in parentheses () are coefficients of $p_i p_j p_k$

terms in braces { } are coefficients of $p_i^2 p_j \delta(j,k)$ plus two cyclic permutations

$$[-1,-1,-2,0]_3 = (1/2 * \log u - 1/(u-1)^3 * \log u - 3/2/(u-1)^2 * \log u + 1/(u-1)^2 + 1/(u-1)) \\ + \{ - 1/4 + 1/4 * \log u + 1/4/(u-1)^2 * \log u + 1/2/(u-1) * \log u - u/4 - 1/4/(u-1) \}$$

$$[-1,-2,-1,0]_3 = (1/2 + 1/(u-1)^3 * \log u + 1/(u-1)^2 * \log u - 1/(u-1)^2 - 1/2/(u-1)) \\ + \{ 1/4 - 1/4 * \log u - 1/4/(u-1)^2 * \log u - 1/2/(u-1) * \log u + 1/4/(u-1) \}$$

$$[-1,-1,-2,1]_3 = (- 1/6 + 1/3 * \log u * (1 - 1/(u-1))^3 - 1/(u-1)^2 + 1/(u-1)) + 1/3/(u-1)^2 \\ + 1/6/(u-1)) + \{ \log u + 1/(u-1)^2 * \log u + 2/(u-1) * \log u - u - u/(u-1) \}/12$$

$$[-1,-2,-1,1]_3 = (1/3 + 1/3/(u-1)^3 * \log u + 1/6/(u-1)^2 * \log u - 1/6/(u-1) * \log u - 1/3/(u-1)^2) \\ + \{ 1/12 - 1/12 * \log u - 1/12/(u-1)^2 * \log u - 1/6/(u-1) * \log u + 1/12/(u-1) \}$$

$$[-1,-1,-2,2]_3 = (- 1/6 + \log u * (1/4 - 1/6/(u-1))^3 - 1/12/(u-1)^2 + 1/3/(u-1)) + 1/6/(u-1)^2 \\ + \{ 1/24 * \log u + 1/24/(u-1)^2 * \log u + 1/12/(u-1) * \log u - u/24 - u/24/(u-1) \}$$

$$[-1,-2,-1,2]_3 = (1/4 + 1/6/(u-1)^3 * \log u - 1/6/(u-1) * \log u - 1/6/(u-1)^2 + 1/12/(u-1)) \\ + \{ - 1/24 * \log u - 1/24/(u-1)^2 * \log u - 1/12/(u-1) * \log u + u/24/(u-1) \}$$

TABLE 8(B) $K = -1$, $\alpha_1 = -1$

$$[K,M,N,L]_3 = \int d^4q ((p-q)^2)^K (q^2)^M (q^+)^N (q^-)^L q_i q_j q_k$$

terms in parentheses () are coefficients of $p_i p_j p_k$

terms in braces {} are coefficients of $p_i^2 \delta(j,k)$ plus two cyclic permutations

$$[-1,0,-2,0]_3 = (- 3/2 - u/2 - 3/2/(u-1)) + \{ u^2/4 \log u + 3u^2/8 \} + \{ 1/\hat{e}_1 * (u^2/4) \}$$

$$[-1,-1,-1,0]_3 = (- 1/2 + 1/3 \log u * (1 + 1/(u-1)^3) - 1/3/(u-1)^2 - 5/6/(u-1)) + \{ u/6 \hat{e}_1 \} \\ + \{ 1/12 \log u + u/6 \log u - 1/12/(u-1)^2 \log u + u/18 + u/12/(u-1) \}$$

$$[-1,-2,0,0]_3 = (1/3 - 1/2/(u-1)) + \{ 1/36 \} + 1/12 * \{ 1/\hat{e}_1 \}$$

$$[-1,0,-2,1]_3 = (- 1/2 - 1/2/(u-1)) + \{ u^2/12 \log u + 7u^2/72 \} + \{ 1/\hat{e}_1 * (u^2/12) \}$$

$$[-1,-1,-1,1]_3 = (- 1/6 + 1/4 \log u + 1/12/(u-1)^3 \log u - 1/12/(u-1)^2 \log u + 1/12/(u-1) \log u \\ - 1/12/(u-1)^2 - 1/4/(u-1)) + \{ 1/\hat{e}_1 * (u/16) \} \\ + \{ 1/24 \log u + u/16 \log u - 1/48/(u-1)^2 \log u + 1/48/(u-1) \log u + u/48/(u-1) \}$$

$$[-1,-2,0,1]_3 = (1/4 - 1/4/(u-1)) + \{ 1/72 \} + 1/24 * \{ 1/\hat{e}_1 \}$$

$$[-1,0,-2,2]_3 = (- 1/4 + u/12 - 1/4/(u-1)) + \{ u^2/24 \log u + 11u^2/288 \} + \{ 1/\hat{e}_1 * (u^2/24) \}$$

$$[-1,-1,-1,2]_3 = (- 1/12 + 1/5 \log u + 1/30/(u-1)^3 \log u - 1/15/(u-1)^2 \log u + 1/10/(u-1) \log u \\ - 1/30/(u-1)^2 - 7/60/(u-1)) + \{ 1/\hat{e}_1 * (u/30) \} \\ + \{ u/120/(u-1) + \log u * (1/40 + u/30 - 1/120/(u-1)^2 + 1/60/(u-1)) - 7u/900 \}$$

$$[-1,-2,0,2]_3 = (1/5 - 3/20/(u-1)) + \{ 1/150 \} + 1/40 * \{ 1/\hat{e}_1 \}$$

TABLE 8(C) $K = -1$, $\alpha_1 = 0$

$$[K,M,N,L]_3 = \int d^4q ((p-q)^2)^K (q^2)^M (q^+)^N (q^-)^L q_1 q_j q_k$$

terms in parentheses () are coefficients of $p_i p_j p_k$

terms in braces { } are coefficients of $p_i^2 \delta(j,k)$ plus two cyclic permutations

$$\begin{aligned} [-1,1,-2,0]_3 &= (7/2 - 5u^2/2 \log u + 9u/2 + 21u^2/4 + 7/2/(u-1)) \\ &+ (1/\hat{e}_1 * (- 5u^2/2)) + \{ 1/\hat{e}_1 * (u^2/4 - 5u^3/6) \} \\ &+ \{ u^2/4 \log u - 5u^3/6 \log u + 3u^2/8 - 8u^3/9 \} \end{aligned}$$

$$\begin{aligned} [-1,0,-1,0]_3 &= (3/2 - u \log u + 5u/2 + 3/2/(u-1)) + (1/\hat{e}_1 * (- u)) \\ &+ \{ - u^2/4 \log u - u^2/8 \} + \{ 1/\hat{e}_1 * (- u^2/4) \} \end{aligned}$$

$$[-1,-1,0,0]_3 = (7/12 + 1/4/(u-1)) - (1/4 * 1/\hat{e}_1) + \{1/36\} - 1/24 * \{ 1/\hat{e}_1 \}$$

$$[-1,-2,1,0]_3 = (1/4 - 1/4/(u-1)) + \{1/72\} + 1/24 * \{ 1/\hat{e}_1 \}$$

$$\begin{aligned} [-1,1,-2,1]_3 &= (1 - u^2 \log u + 3u/2 + 2u^2 + 1/(u-1)) + (1/\hat{e}_1 * (- u^2)) \\ &+ \{ u^2/12 \log u - u^3/4 \log u + 7u^2/72 - 3u^3/16 \} \\ &+ \{ 1/\hat{e}_1 * (u^2/12 - u^3/4) \} \end{aligned}$$

$$\begin{aligned} [-1,0,-1,1]_3 &= (1/2 - u/2 \log u + 5u/4 + 1/2/(u-1)) + (1/\hat{e}_1 * (- u/2)) \\ &+ \{ - u^2/12 \log u - u^2/72 \} + \{ 1/\hat{e}_1 * (- u^2/12) \} \end{aligned}$$

$$[-1,-1,0,1]_3 = (149/300 + 3/20/(u-1)) - (1/5 * 1/\hat{e}_1) + \{11/600\} - 1/40 * \{ 1/\hat{e}_1 \}$$

$$\begin{aligned} [-1,-2,1,1]_3 &= (1/5 - 1/75/u - 1/20/(u-1)) + (1/\hat{e}_1 * (- 1/20/u)) \\ &+ \{ 1/150 + 17/1800/u \} + \{ 1/\hat{e}_1 * (1/40 - 1/60/u) \} \end{aligned}$$

$$\begin{aligned} [-1,1,-2,2]_3 &= (9/20 - 7u^2/12 \log u + 47u/60 + 839u^2/720 + 9/20/(u-1)) \\ &+ (1/\hat{e}_1 * (- 7u^2/12)) + \{ 1/\hat{e}_1 * (u^2/24 - 7u^3/60) \} \\ &+ \{ u^2/24 \log u - 7u^3/60 \log u + 11u^2/288 - 211u^3/3600 \} \end{aligned}$$

$$[-1,0,-1,2]_3 = (1/4 - u/3 \cdot \log u + 31u/36 + 1/4/(u-1)) + (1/\hat{e}_1 * (- u/3)) \\ + \{ - u^2/24 \cdot \log u + u^2/288 \} + \{ 1/\hat{e}_1 * (- u^2/24) \}$$

$$[-1,-1,0,2]_3 = (157/360 + 1/10/(u-1)) - (1/6 \cdot 1/\hat{e}_1) + \{49/3600\} - 1/60 \cdot \{ 1/\hat{e}_1 \}$$

$$[-1,-2,1,2]_3 = (1/6 + 19/900/u) + (1/\hat{e}_1 * (- 1/15/u)) \\ + \{ 11/3600 + 17/1800/u \} + \{ 1/\hat{e}_1 * (1/60 - 1/60/u) \}$$

TABLE 8(D) $K = -1$, $\alpha_1 = 1$

$$[K,M,N,L]_3 = f \int d^4 q ((p-q)^2)^K (q^2)^M (q^+)^N (q^-)^L q_i q_j q_k$$

terms in parentheses () are coefficients of $p_i p_j p_k$

terms in braces { } are coefficients of $p_i^2 p_j \delta(j,k)$ plus two cyclic permutations

$$\begin{aligned} [-1,2,-2,0]_3 &= (-21/4 - 6u^2 \log u + 14u^3 \log u - 17u/4 - 11u^2/4 - 103u^3/4 - 21/4/(u-1)) \\ &+ (1/\hat{e}_1 * (-6u^2 + 14u^3)) + \{ 1/\hat{e}_1 * (u^2/4 - 2u^3 + 21u^4/8) \} \\ &+ \{ u^2/4 \log u - 2u^3 \log u + 21u^4/8 \log u + 3u^2/8 - 2u^3 + 73u^4/32 \} \end{aligned}$$

$$\begin{aligned} [-1,1,-1,0]_3 &= (-3/2 - u \log u + 3u^2 \log u - u/2 - 13u^2/2 - 3/2/(u-1)) \\ &+ (1/\hat{e}_1 * (-u + 3u^2)) + \{ 1/\hat{e}_1 * (-u^2/4 + u^3/2) \} \\ &+ \{ -u^2/4 \log u + u^3/2 \log u - u^2/8 + u^3/4 \} \end{aligned}$$

$$[-1,0,0,0]_3 = 0$$

$$[-1,-1,1,0]_3 = (149/300 + 3/20/(u-1)) - (1/5 * 1/\hat{e}_1) + \{ 11/600 \} - 1/40 * \{ 1/\hat{e}_1 \}$$

$$[-1,-2,2,0]_3 = (1/5 - 3/20/(u-1)) + \{ 1/150 \} + 1/40 * \{ 1/\hat{e}_1 \}$$

$$\begin{aligned} [-1,2,-2,1]_3 &= (-6/5 - 7u^2/3 \log u + 14u^3/3 \log u - 7u/10 + 26u^2/45 - 409u^3/45 - 6/5/(u-1)) \\ &+ (1/\hat{e}_1 * (-7u^2/3 + 14u^3/3)) + \{ 1/\hat{e}_1 * (u^2/12 - 7u^3/12 + 7u^4/10) \} \\ &+ \{ u^2/12 \log u - 7u^3/12 \log u + 7u^4/10 \log u + 7u^2/72 - 59u^3/144 + 41u^4/100 \} \end{aligned}$$

$$\begin{aligned} [-1,1,-1,1]_3 &= (-3/8 - u/2 \log u + 7u^2/6 \log u + 3u/8 - 193u^2/72 - 3/8/(u-1)) \\ &+ (1/\hat{e}_1 * (-u/2 + 7u^2/6)) + \{ 1/\hat{e}_1 * (-u^2/12 + 7u^3/48) \} \\ &+ \{ -u^2/12 \log u + 7u^3/48 \log u - u^2/72 + 17u^3/576 \} \end{aligned}$$

$$[-1,0,0,1]_3 = 0$$

$$\begin{aligned} [-1,-1,1,1]_3 &= (157/360 - 31/450/u + 3/40/(u-1)) + \{ 49/3600 - 1/225/u \} \\ &+ (1/\hat{e}_1 * (-1/6 + 1/30/u)) + \{ 1/\hat{e}_1 * (-1/60 + 1/240/u) \} \end{aligned}$$

$$\begin{aligned} [-1,-2,2,1]_3 &= (1/6 + 19/900/u) + \{ 11/3600 + 17/1800/u \} \\ &+ (1/\hat{e}_1 * (-1/15/u)) + \{ 1/\hat{e}_1 * (1/60 - 1/60/u) \} \end{aligned}$$

$$\begin{aligned}
[-1,2,-2,2]_3 &= (-9/20 - 4u^2/3 \log u + 12u^3/5 \log u - 7u/60 + 41u^2/45 - 371u^3/75 - 9/20/(u-1)) \\
&+ (1/\hat{e}_1 * (-4u^2/3 + 12u^3/5)) + \{ 1/\hat{e}_1 * (u^2/24 - 4u^3/15 + 3u^4/10) \} \\
&+ \{ u^2/24 \log u - 4u^3/15 \log u + 3u^4/10 \log u + 11u^2/288 - 28u^3/225 + 8u^4/75 \}
\end{aligned}$$

$$\begin{aligned}
[-1,1,-1,2]_3 &= (-3/20 - u/3 \log u + 2u^2/3 \log u + 83u/180 - 73u^2/45 - 3/20/(u-1)) \\
&+ (1/\hat{e}_1 * (-u/3 + 2u^2/3)) + \{ 1/\hat{e}_1 * (-u^2/24 + u^3/15) \} \\
&+ \{ -u^2/24 \log u + u^3/15 \log u + u^2/288 - u^3/450 \}
\end{aligned}$$

$$[-1,0,0,2]_3 = 0$$

$$\begin{aligned}
[-1,-1,1,2]_3 &= (383/980 - 967/8820/u + 3/70/(u-1)) \\
&+ (1/\hat{e}_1 * (-1/7 + 1/21/u)) + \{ 1/\hat{e}_1 * (-1/84 + 1/210/u) \} \\
&+ \{ 379/35280 - 463/88200/u \}
\end{aligned}$$

$$\begin{aligned}
[-1,-2,2,2]_3 &= (1/7 - 1241/44100/u^2 + 779/8820/u + 3/140/(u-1)) \\
&+ (1/\hat{e}_1 * (2/105/u^2 - 2/21/u)) + \{ 1/\hat{e}_1 * (1/84 + 1/280/u^2 - 2/105/u) \} \\
&+ \{ 41/35280 - 107/29400/u^2 + 253/22050/u \}
\end{aligned}$$

TABLE 8(E) $K = -1$, $\alpha_1 = 2$

$$[K,M,N,L]_3 = f \int d^4q ((p-q)^2)^K (q^2)^M (q^+)^N (q^-)^L q_i q_j q_k$$

terms in parentheses () are coefficients of $p_i p_j p_k$

terms in braces { } are coefficients of $p_i^2 \delta(j,k)$ plus two cyclic permutations

$$\begin{aligned} [-1,2,-1,0]_3 &= (-u \log u + 7u^2 \log u - 28u^3/3 \log u + 5u/2 - 17u^2/2 + 172u^3/9 + 3u/2/(u-1)) \\ &+ (1/\hat{e}_1 * (-u + 7u^2 - 28u^3/3)) + \{ 1/\hat{e}_1 * (-u^2/4 + 7u^3/6 - 7u^4/6) \} \\ &+ \{ -u^2/4 \log u + 7u^3/6 \log u - 7u^4/6 \log u - u^2/8 + 19u^3/36 - 19u^4/36 \} \\ [-1,1,0,0]_3 &= 0 \\ [-1,0,1,0]_3 &= 0 \\ [-1,-1,2,0]_3 &= (157/360 + 1/10/(u-1)) - (1/6 * 1/\hat{e}_1) + \{ 49/3600 \} - 1/60 * \{ 1/\hat{e}_1 \} \\ [-1,2,-1,1]_3 &= (3u/10/(u-1) - \log u * (u/2 - 8u^2/3 + 3u^3) + 21u/20 - 199u^2/45 + 199u^3/30) \\ &+ (1/\hat{e}_1 * (-u/2 + 8u^2/3 - 3u^3)) + \{ 1/\hat{e}_1 * (-u^2/12 + u^3/3 - 3u^4/10) \} \\ &+ \{ -u^2/12 \log u + u^3/3 \log u - 3u^4/10 \log u - u^2/72 + u^3/18 - 17u^4/300 \} \\ [-1,1,0,1]_3 &= 0 \\ [-1,0,1,1]_3 &= 0 \\ [-1,-1,2,1]_3 &= (383/980 - 967/8820/u + 3/70/(u-1)) + (1/\hat{e}_1 * (-1/7 + 1/21/u)) \\ &+ \{ 379/35280 - 463/88200/u \} + \{ 1/\hat{e}_1 * (-1/84 + 1/210/u) \} \\ [-1,2,-1,2]_3 &= (u/10/(u-1) - \log u * (u/3 - 3u^2/2 + 3u^3/2) + 32u/45 - 353u^2/120 + 85u^3/24) \\ &+ (1/\hat{e}_1 * (-u/3 + 3u^2/2 - 3u^3/2)) + \{ 1/\hat{e}_1 * (-u^2/24 + 3u^3/20 - u^4/8) \} \\ &+ \{ -u^2/24 \log u + 3u^3/20 \log u - u^4/8 \log u + u^2/288 - 11u^3/1200 + u^4/288 \} \\ [-1,1,0,2]_3 &= 0 \\ [-1,0,1,2]_3 &= 0 \\ [-1,-1,2,2]_3 &= (199/560 + 67/4410/u^2 - 71/392/u + 3/140/(u-1)) + \{ 69/7840 + 71/88200/u^2 - 121/17640/u \} \\ &- (1/\hat{e}_1 * (1/8 + 1/168/u^2 - 1/14/u)) - \{ 1/\hat{e}_1 * (1/112 + 1/1680/u^2 - 1/168/u) \} \end{aligned}$$

ISSN 0067 - 0367

To identify individual documents in the series
we have assigned an AECL- number to each.

Please refer to the AECL- number when re-
questing additional copies of this document

from

Scientific Document Distribution Office
Atomic Energy of Canada Limited
Chalk River, Ontario, Canada
K0J 1J0

Price: B

ISSN 0067 - 0367

Pour identifier les rapports individuels faisant
partie de cette série nous avons assigné
un numéro AECL- à chacun.

Veuillez faire mention du numéro AECL- si
vous demandez d'autres exemplaires de ce
rapport

au

Service de Distribution des Documents Officiels
L'Energie Atomique du Canada Limitée
Chalk River, Ontario, Canada
K0J 1J0

Prix: B

©ATOMIC ENERGY OF CANADA LIMITED, 1987